

EMBARGOED UNTIL March 2, 2009 at 0900 CET



HUNGRY OCEANS: WHAT HAPPENS WHEN THE PREY IS GONE?

Margot L. Stiles, Laure Katz, Tess Geers, Sarah Winter, Ellycia Harrould-Kolieb, Andrew Collier, Ben Enticknap,
E. Kate Barnes, Sarah Hale, Prisca Faure, Jaroslava Waters, Michael F. Hirshfield

oceana.org/prey

Acknowledgements

Bryant Austin
Rob Birnbaum
Dee Boersma
Andre Boustany
Sam Breach
Pablo Andrés Cáceres Contreras
Cathou Cathare
Eric Cheng
John Croxall
Chris Dent
Wetjens Dimmlich
FishBase
Pam Lyons Gromen, National Center for Marine Conservation
Anne Guerry
Roger T. Hanlon
Gene Helfman
Jennifer Jacquet
Kev Jacobs
Shannon Johnson
Steve Johnson
Uwe Kils
Marine Fish Conservation Network
Christopher Michel, fotografica.us
Paul McDonnell
Alexander Perry, Northumbria Photography
Jim Price, Chesapeake Bay Ecological Foundation
John Rix, Fathom This Underwater Productions
Colin Robson
Horeal Vidal Sabatte
Aziz T. Saltik
Gordon Stroupe


Oceana staff including Jim Ayers, Gib Brogan, Alejandro Buschmann, María José Cornax, Keith Ellenbogen, Suzannah Evans, Cristián Gutiérrez, Chris Krenz, Patricia Lastra, Jesús Renedo, Santi Roberts, Nausheen Saeed, Jon Warrenchuk

Design by Arista Advertising
Cover Photo by Colin Robson

Images in this report may not be reproduced without permission. Information in this report may be reproduced as long as appropriately credited. Line drawings throughout the report are from the Alaska Department of Fish and Game, ASMFC, FAO, Ocean Institute, UC Berkeley, and the Virginia Institute of Marine Science.

TABLE OF CONTENTS

2	Overview Predators Need Their Prey Ecosystem Resilience
4	Predators Bluefin Tuna and Other Big Fish Whales, Penguins, and Other Animals
13	Overexploited Prey Species
15	Threats to Prey Species Overfishing Aquaculture Climate Change
23	Solutions Getting Started Looking Ahead Solutions Summary
28	Prey Species Around the World
31	Resources





Overview

According to conventional wisdom, small, fast-growing fish are impossible to overfish because their populations are so large and grow so quickly. Yet we are now seeing disquieting signs that conventional wisdom is wrong. Most significantly, scientists are reporting ocean predators emaciated from lack of food, vulnerable to disease and without enough energy to reproduce. Scrawny predators—dolphins, striped bass, and even whales—have turned up along coastlines around the world. Recreational fishermen are losing both their target fish—and their bait. Fishing communities are losing their livelihoods.


Because we have overlooked hungry predators, we have allowed overfishing of their prey, causing food shortages in the ocean. In addition, by fishing where and when they are breeding, we are driving prey populations to the brink of disaster, and in some cases beyond it.

At the same time, our continued demand for salmon, tuna, and other large predators has driven explosive growth in aquaculture. Rather than relieving pressure on wild fish, growing these large carnivores requires a steady supply of prey that are caught and ground into oil and meal. As the industry grows, it is straining the existing supply of prey fish, putting additional pressure on populations that cannot supply the demand.

Because populations of many small prey species are sensitive to changes in temperature, ocean currents, and El Niño, they are particularly vulnerable to climate change. For predators, even small climate-driven shifts in the local availability of squid and other prey during breeding can lead to malnourished young or abandonment.

For many prey species, humans have replaced their natural predators—the other fish, the sea birds, the marine mammals—all the species that depend on them for their existence. Fishermen who catch prey species are beginning to turn up empty nets. Unless the current trends are reversed, we can look forward to a future with increasingly hungry oceans.

Colin Robson



Prey species
underpin
marine food
webs around
the world

*Seabirds dive
on a herring
school*

PREDATORS NEED THEIR PREY

The great predators of the ocean spend most of each day hunting for food. Scientists studying sperm whales estimate they spend nearly three quarters of their time searching for squid and other prey (Watwood et al. 2006). Abundant schooling fish fuel the blue marlin's speed and strength (Abitia Cardenas 1999), and are staples in the diet of many whales.

Hungry animals may fail to nurse or find enough food for their young, and sometimes skip breeding season entirely. During an eight-year prey shortage in the Faroe Islands, no Arctic tern chicks survived (Wright et al. 1996). During another food shortage, Galápagos penguins were forced to "desert their eggs and chicks to search for food to save themselves while their chicks starved to death." (Boersma et al. 2008) When long-lived animals like whales go hungry, the next generation is at risk.

Predators are normally forced to rely on less desirable, less nutritious, or less abundant prey for short periods of time. If this happens too often or for long periods of time, however, predators can become malnourished and vulnerable to disease. Poor health and food shortages may have left striped dolphins vulnerable to a 1990s plague in the Mediterranean that led to many deaths (Aguilar and Raga in Bearzi et al. 2003).

ECOSYSTEM RESILIENCE AT RISK

Only now that predators are going hungry are forage fish becoming recognized for their role at the foundation of marine food webs. Predators consume great quantities of tiny fish—often all the same species. Within an ecosystem perhaps only two or three species fill that role (Cury et al. 2000). If one prey population crashes, few options are left for its predators. This low level of redundancy can result in a lack of resilience to other stresses on the ecosystem.

Shannon Johnson





PREDATORS:

Bluefin Tuna and Other Big Fish

Loss of prey for large predatory fish translates to loss of prey for commercial fishermen and recreational anglers, as tuna, salmon, and striped bass go hungry. Species in recovery such as North Atlantic Bluefin Tuna or Striped Bass are particularly in need of abundant prey to rebuild their populations from overfishing.

Northern Atlantic Bluefin Tuna

Bluefin tuna (*Thunnus thynnus*) are some of the largest and fastest fish in the oceans—and also the most valuable, with a record of \$173,600 for a single fish (Associated Press 2001). This demand has driven worldwide overfishing of this species, and in the western Atlantic populations have been reduced by more than 80 percent since 1975 (ICCAT 2006). Surprisingly, the basic biology of this fish is just beginning to be understood, including its food-centric migration.

Though bluefin tuna are top predators and opportunistic feeders, their diet is often dominated by one or two favorite prey species that provide optimal sources of energy (Chase 2002). Mass movements of bluefin are synchronized with spawning and feeding schools of various species of prey fish off the east coast of North America. At each step along the way, changing prey populations substantially affect regional aggregations of bluefin tuna, as illustrated in Figure 1.




Figure 1. Seasonal prey of Atlantic Bluefin Tuna off the Northeast US
 Source: Data in percent of bluefin tuna diet by weight from Chase 2002

Bluefin tuna in the Mediterranean



Oceana / Keith Ellenbogen

During winter months Atlantic bluefin tuna concentrate off the coast of North Carolina where they feast on Atlantic menhaden (*Brevoortia tyrannus*), the staple of their diet. In a recent study of stomach contents from commercially caught tuna, Atlantic menhaden were found 85 percent of the time and made up 95 percent of the bluefin prey by weight (Butler 2007). While predators such as bluefish, striped bass, and weakfish (as well as the commercial fishery) consume much greater quantities of menhaden than tuna, menhaden are essential to bluefin on their winter foraging grounds.

Bluefin tuna concentrate along the southeastern coast of the US during fall and winter, moving north through spring and summer to rich feeding grounds of the North Atlantic (Block et al. 2005). At each location their diet is dominated by one or two preferred fish or squid (Chase 2002). When Atlantic menhaden disperse from concentrated aggregations in North Carolina towards the end of March, bluefin tuna move offshore. Here their diet is poorly known, but they may shift to a wider variety of fish and squid (Dragovich 1970, Boustany, pers. comm.). In June and July bluefin tuna return inshore to New England feeding grounds and a diet dominated by one or two major prey species (Chase 2002).

Mediterranean bluefin tuna feed on squid and small fish, depending on where they live. For juvenile tuna, small schooling fish are particularly important (Sarà and Sarà 2007, Sinopoli et al. 2004).

Pacific Salmon

Like the bluefin tuna, wild Pacific salmon (*Chinook *Oncorhynchus tshawytscha*, Coho *Oncorhynchus kisutch*) prey on krill and small fish throughout their migrations. As they leave their home rivers for the open sea, young chinook and coho salmon (*Oncorhynchus spp.*) feed heavily on Pacific herring, Pacific sand lance, and surf smelt. These prey species spawn in the intertidal zone of many beaches along the Pacific Northwest coast and serve as the first meal for juvenile salmon as they reach the sea, making possible their journey and survival into adulthood. Full-grown salmon range along the west coast of the United States, shifting between springtime krill, crabs and squid to a summertime diet of anchovy, smelt, and sand lance. (FishBase, Hunt 1999, Tyler et al. 2001, Zavolokin et al. 2007, Sakai et al. 2005, Sagawa et al 2007)*

Chinook or King Salmon change their diet seasonally



Save Our Wild Salmon

oceana.org/prey



Striped Bass

Striped bass (*Morone saxatilis*) also track the movements of their most important prey, migrating along the coast with Atlantic menhaden and moving between salt and fresh water with river herring. Each spring when striped bass are concentrated in the Chesapeake and Delaware Bay, their diet is dominated by Atlantic menhaden and river herring, and their diets shift to sand lance (also known as sand eel) and other prey as they migrate north each summer (Walter et al. 2003). In fall and winter, striped bass return to focus on menhaden in the bays of Maryland, Delaware, and North Carolina.

Striped bass may struggle with the steady decline of their major prey item, Atlantic menhaden (Uphoff 2003, D.Russell and J.Price, pers. comm.). Menhaden, herring, and bay anchovy are all important prey (Griffin and Magruff 2003, Manooch 1973).

TABLE 1. Recreational and Commercial Fishery Species Dependent on Prey

TARGET SPECIES	STATUS (FAO AND OTHER SOURCES)	SELECT PREY SPECIES
Pacific Salmon	Fully exploited to overexploited	Krill, Squid, Sand Lance, Herring, Sardine, Northern Anchovy, Juvenile Rockfish, Juvenile Atka Mackerel, Walleye Pollock, Lanternfish
Bluefin Tuna	Depleted to severely overexploited	Menhaden, Sand Lance, Herring, Mackerel, Squid
Striped Bass	Recovered	Menhaden, Bay Anchovy, Herring
Pacific Halibut	Fully exploited	Sand Lance, Herring, Walleye Pollock, Squid
South Pacific Hake	Fully exploited to depleted	Anchovy, Anchoveta, Squid, Sardine, Krill
European Hake	Overexploited	Blue Whiting, European Anchovy, European Pilchard, Lanternfish, Horse Mackerel
Southern Hake	Fully exploited to overexploited	Squid, Krill, Blue Whiting

Sources: Butler 2007, Boustany pers. comm., Cartes et al. 2004, Chase 2002, Dragovich 1970, Fishbase, Hunt 1999, IPHC 1998, FAO 2005, Mahe et al. 2007, Paya 1992, Sagawa et al. 2007, Sakai et al. 2005, Tam et al. 2006, Tyler et al. 2001, Velasco 1998, Walter et al. 2003, Zavolokin et al. 2007



John Rix / Fathom This Underwater Productions



Save Our Wild Salmon



Beccy Breach

“One cannot think well, love well,
sleep well, if one has not dined well.”

— Virginia Woolf



A humpback whale calf swims ahead of its mother

Bryant Austin

PREDATORS: Whales, Penguins and Other Animals

Ocean predators also include whales, dolphins, seals, sea lions, and birds. Although many of these animals are protected under national and international law, they remain vulnerable to food shortages. They depend on nearby access to squid, krill, and small fish to provide energy reserves for daily survival and for their young.

Endangered species are especially sensitive to food shortages and need abundant food to rebuild their populations. Starving whales are unable to nurse their young, and hungry seabirds may become susceptible to disease. Most animals feed heavily to gain weight before giving birth or after migration, and abundant prey is critical at these times.


Blue Whales in Southern Chile

The largest animals that have ever lived on earth are powered entirely by krill, and blue whales (*Balaenoptera musculus*) eat a lot to maintain their considerable bulk at 100 feet (30 m) and 200 tons (180 mt) (Clapham et al. 1999, Hucke-Gaete et al. 2006). Blue whales were hunted to the brink of extinction in the 20th century, reduced in the Southern Hemisphere to less than 3 percent of their original numbers (Hucke-Gaete 2003). Now blue whales need more food than ever, to fuel reproduction so their populations can recover.

Krill hotspots such as the Chiloé-Corcovado region in Southern Chile have historically been recognized as blue whale feeding sites. In 1907 a Norwegian whaling ship caught 37 whales in the area and in recent years blue whales have been sighted with increasing frequency; groups of more than 60 whales and cow-calf pairs have been observed (Hucke-Gaete et al. 2006, Hucke-Gaete 2003). The main reason why Southern Hemisphere blue whale populations nurse their calves in Chiloé-Corcovado is to take advantage of the abundant krill supply. This tiny crustacean fuels one of the most important feeding and nursing grounds for Southern Hemisphere blue whales, and is responsible for their gradual recovery from whaling.

Antarctic krill
form large swarms





Dolphins in the Mediterranean

Mediterranean bottlenose dolphins (*Tursiops truncatus*) have declined significantly, in part because of overfishing of sardines and anchovy that also led to collapse of the fishery in 1987 (Politi 2000). Scientists working in the eastern Ionian Sea found 40 percent of bottlenose dolphins visibly emaciated due to starvation and other causes (Politi et al. 2000).

Short-beaked common dolphins (*Delphinus delphis*) were formerly abundant throughout the Mediterranean, but are now concentrated in a much narrower range near Algeria, Tunisia, Malta, and in the Alboràn, Aegean, Tyrrhenian, and eastern Ionian Seas. Several factors likely contributed to their decline, including reduced access to prey due to overfishing (Bearzi et al. 2003). Poor nutrition may also increase dolphins' susceptibility to disease, an important factor in two mass mortality events in the Mediterranean (Bearzi et al. 2003).



Jesús Renedo / Oceana

A dolphin leaps from the Mediterranean

Marbled Murrelet

Deforestation, nest predation, and oil spills have previously been blamed for driving the marbled murrelet (*Brachyramphus marmoratus*) of central California to the endangered species list. While these factors clearly contributed, recent evidence suggests that the overfishing of sardines and other prey species is also partly to blame. The diet of the marbled murrelet in the Monterey Bay ecosystem has drastically been altered over the past century (Becker and Beissinger 2006). Fisheries declines, especially the infamous 1950s collapse of the California sardine fishery, have reduced the availability of fish as food for the marbled murrelet. To make matters worse, a marbled murrelet must spend added time and energy catching 80 krill to match the energy found in a single Pacific sardine (Becker and Beissinger 2006).



Magellanic Penguin

Penguins, cormorants, terns and other bird species are currently threatened by a developing anchovy fishery in the Patagonian ecosystem (Skegwar et al. 2007). The Southwest Atlantic anchovy (*Engraulis anchoita*) compose more than half of the Magellanic penguin diet (Skegwar 2007). Despite the central importance of anchovy to the Patagonian food web and the natural variation in its availability, some have pushed to expand the anchovy fishery. If the needs of the ecosystem as a whole are ignored, this fishery could drastically change the food web, with dire consequences for the seabirds of Patagonia.

TABLE 2. Endangered and Protected Species Dependent on Prey Species

ENDANGERED OR PROTECTED SPECIES	STATUS (IUCN)	SELECT PREY SPECIES
Sperm whale	Vulnerable	Squid
Blue whale	Endangered	Krill
Dolphins in the Mediterranean	Data deficient for some species	Anchovy, Sardine, Hake, Whiting, Cephalopods
Harbor porpoise	Vulnerable	Whiting, Herring, Sand Lance, Capelin, Cephalopods
Steller sea lion	Endangered	Sand Lance, Pollock, Herring, Capelin, Squid
Magellanic penguin	Near threatened	Anchovy, Cuttlefish, Hake, Squid, Krill
Marbled murrelet	Endangered	Sardine, Anchovy, Squid, Sand Lance, Herring, Krill, Capelin
Kittiwake	Red-legged: Vulnerable Black-legged: Least concern	Pollock, Herring, Sand Lance, Capelin
Puffin	Least concern	Capelin, Herring, Sand Lance, Squid, Lanternfish

Sources: Baillie and Jones 2003, Barrett and Furness 1990, Bearzi et al. 2003, Blanco et al. 2001, Börjesson 2002, Burkett 1995, COSEWIC 2003, Durant 2003, Falk 1992, Forero et al. 2002, Gandini et al. 1999, Gellatt et al. 2007, Huckle-Gaete et al. 2006, MacLeod et al. 2007, Ozturk et al. 2007, Pinto et al. 2006, Raga et al. 2006, Rodway and Montevecchi 1996, Santos and Pierce 2003, Sanger 1987, Scolaro et al. 1999, Silva 1999, Springer 1986, Spitz et al. 2007, Suryan 2000, Tonay 2007, Trites et al. 2007, Vanda et al. 2001, Wanless et al. 2004, Watwood et al. 2006, Winship and Trites 2003




Weijens Dimmlich

Freshly caught sardines

OVEREXPLOITED PREY SPECIES

Fisheries targeting prey species have grown dramatically during the past century and are currently overdrawn, leaving predators with depleted food supplies. Early fisheries targeted only herring, sardines, and menhaden until expanding to meet the demand for cheap animal feed after World War II, and the more recent demand from carnivorous aquaculture (IFFO 2006a, Alder and Pauly 2006, Watson et al. 2006). Now more than 88 different prey stocks are caught everywhere from the tropics to the poles (FAO).

Ten Biggest Fisheries in the World



Prey species have become targets for the largest fisheries in the world, as populations of the bigger fish in the sea are exhausted (Jackson 2008, Pauly et al. 1998). Seven of the top ten fisheries rely on prey fish, as illustrated in Figure 2, and today's landings of prey fish are more than four times those of 1950 (FAO). More than 10 million metric tons of anchoveta alone are removed from the ocean every year, made unavailable as prey to seabirds, mammals, and predatory fish.

FIGURE 2. Ten Biggest Fisheries in the World
Source: FAO 2006

Squid fishing

ANONYMOUS FISH

Prey are known by many names, in part because they are usually overlooked. Some are accurate yet vague, including “small pelagics” indicating that they are found in open water, or “schooling fish.” Others such as “bait fish,” “forage fish,” and “prey fish” define them by their uses and fail to include non-fish species like squid and krill.



Horeal Vidal Sabatte

oceans.org/prey




FIGURE 3. Prey Stocks that are Fully or Overfished
Source: Percentage of 88 assessed stocks from FAO 2005

Decades of intense fishing pressure have taken their toll. Among the stocks assessed by the FAO, the majority are fully or overexploited with no regard for predator needs (2005). For an additional 16 percent, their status is unknown and yet fishing continues. This leaves only 20 percent of currently fished prey populations with any potential to sustain natural predators (see Figure 3). Even fewer of these populations remain large enough to support growing pressures from fisheries, aquaculture, and climate change.

According to the United Nations Food and Agricultural Organization, all prey fisheries in the Southeast Pacific are between fully and overexploited, with the exception of squid. The fishery for sardina del Norte (*Sardinops sagax*) peaked in the mid-1980s before crashing to less than one percent of peak landings with no sign of recovery. Nearly all fisheries for prey in the North Atlantic, Central Atlantic, and North Pacific are fully exploited with no consideration for predators. This includes menhaden along the US coast, sand eels in Europe, and California market squid (*Loligo opalescens*).



Rob Birnbaum

Schooling squid

THREATS: OVERFISHING

Prey fish are taken for granted in fisheries management, despite their critical role in marine food webs. Even in planning the recovery of endangered or overfished species, managers often give no consideration of the food supply needed for their populations to rebound. Squid, krill, and other prey remain ignored, unregulated and unaccounted for.

Disrupted Schooling Behavior

Schooling fish and “bait balls” protect individual fish from natural predators, and schools are also formed during spawning to increase the chances of future generations. Schooling prey are so important to tuna, whales, and other long-lived animals that they drive predator migrations, breeding, and nursing.

Unfortunately, schooling prey fish are easy targets for large nets (Alder et al. 2008, Watson et al. 2006) and can be quickly depleted or dispersed. Predators are forced to compete with industrial fishing vessels, and are sometimes captured incidentally while feeding on schools of fish.



Gordon Stroupe

Schooling prey fish surround a Sand Tiger Shark

Prey remain under the radar,
unregulated and unaccounted for.



Gene Helfman

NOAA



Menhaden

Boom and Bust Populations

Many prey species have short life spans. As a result, some prey species are known for dramatic boom and bust cycles from year to year (Alder et al. 2008), often because an entire generation of fish or squid fails to make it to adulthood. Changes in the prevailing currents or temperature may sweep away newly hatched eggs and larvae (Chavez et al. 2003), or major weather events such as hurricanes can wipe out an entire generation of larval fish. In some cases, population crashes can be linked to particular events including oil spills or pollution (Peterson et al. 2003, Paine et al. 1996).

Overfishing can also drive population crashes by removing juveniles before they are old enough to reproduce, or by directly removing spawning adults. Population crashes may also be caused by increased competition or predation by other species in the food web. All of these factors combine over short periods of time to make it extremely difficult to predict prey availability.

Localized Depletion

Many predators depend on their prey to be available at a particular place and time. Depletion of prey populations in one local area can be extremely disruptive to their predators, even if distant populations remain strong (Furness and Tasker 2000). For example, during nesting season, seabirds have a very limited time away from their young to find food and return. If there are not enough prey where and when the birds expect, both chicks and parents are at risk for starvation or death (Hunt and Furness 1996).

Coastal communities and fishermen are also tied to a particular place and season to make their living. In North Carolina and in New England, bluefin tuna fishing depends on tuna arriving during the legal fishing season and staying as long as possible. When the herring is gone, hungry tuna may move on prematurely and the fishermen return empty-handed.

Predators and Natural Mortality

When no information is available on how much ocean predators consume, their needs are arbitrarily assumed to be low. Standard fishery models gloss over the needs of ocean predators in a single number known as “natural mortality” which is often flawed.

For example, recent analysis found that marine mammals and large fish are actually eating four times as many herring than assumed by the official government assessment (Read and Brownstein 2003). In failing to account for these predators, the typical approach overestimates the amount of fishing that can be sustained by their prey. For years we have been catching many more prey fish than can sustain both human fisheries and predators.

Shifting Baselines

Our current management framework assumes ecosystems with very low levels of natural predation. Large predatory fish have been overexploited and the legacy of whaling and depleted marine mammal populations remains. Ironically, as predatory fish recover and reclaim their share of prey species, we blame them for eating too much. In the Chesapeake Bay, some ask whether there will be enough menhaden left as striped bass recover. In the time that striped bass have been absent due to overfishing, human fishing has expanded - removing their prey.

Alexander Perry

Northumbria Photography



A flying puffin

For years we have been fishing on a deficit.



THREATS: AQUACULTURE

Increasingly, the driver behind overfishing of prey species is aquaculture. Salmon, tuna, and other high-value farmed fish are the fastest growing seafood products in the world (Delgado et al. 2003). As a result, an increasing number of new aquaculture operations specialize in fish that eat fish, which require constant supplies of high-calorie feed.

Aquaculture currently consumes more than 81 percent of the prey fish captured and “reduced” to fish oil, and approximately half of those captured for fishmeal (Tacon et al. 2006). The remaining prey fish are used in agriculture and to a lesser degree pet food and pharmaceuticals (Figure 4; Tacon et al. 2006, Delgado et al. 2003, Campbell and Alder 2006, FIN 2007a).

Fishmeal Uses




FIGURE 4. Fishmeal Consumption
Source: Data from Campbell and Alder 2008

Delta Aquaculture Equipment Co.



A third of all global fish landings are destined for fish meal and oil each year, caught mostly in dedicated reduction fisheries (Watson et al. 2006, FIN 2007b, IFFO 2006a, New and Wijkstrom, 2002). These fish are pressed, dried, and milled into concentrated fish oil and brown powder called fishmeal. A smaller percentage of fish meal is derived from byproducts of fish processing (FIN 2007b), and an unknown percentage is derived from unreported or illegal bycatch (Alder et al. 2008, Tuominen and Esmark 2003).

Among farmed fish, salmon consume more fish oil than all other aquaculture operations combined. In 2003, salmon pens alone consumed 51 percent of world fish oil and 19 percent of world fishmeal supplies respectively (FIN 2007b). In fact, the astronomical growth and success of Chile's salmon aquaculture industry is made possible by readily available fish oil supplies produced within the country (Campbell and Alder 2006). For salmon aquaculture, an estimated four to eleven pounds of prey fish are consumed to grow only one pound of farmed salmon (Fishmeal Information Network, Buschmann et al. 2006).

Further growth of the aquaculture industry is limited by the price of aquafeeds (Tuominen and Esmark 2003, Delgado et al. 2003, New and Wijkstrom 2002). Prices for fish oil and meal are expected to increase and will be highest when natural population cycles lead to diminished forage fish catches, creating powerful incentives for overfishing (Delgado et al. 2003).

2020 The year aquaculture outgrows the supply of fishmeal

— New and Wijkstrom, 2002



Eric Cheng

Shoveling prey fish into net pens to feed farmed tuna

TUNA FATTENING

Tuna fattening has expanded in response to continued demand for full-grown bluefin tuna despite reduced supply due to overfishing. Juvenile bluefin are captured and held for varying lengths of time in net pens in Mexico, Canada, Australia, Japan, and throughout the Mediterranean (Ottolenghi et al. 2004, Volpe 2005). They are fed with frozen prey fish and later sold on the international market for sushi and sashimi, particularly in Japan and Korea.

Bluefin tuna are voracious predators, and require between two and ten percent of their body weight in prey fish every day of the peak summer season (Lovatelli 2003). This results in an estimated 225,000 metric tons of prey fish thrown into the Mediterranean tuna pens alone each year (Tudela et al. 2005).

Industry growth is limited by the supply of prey fish, and Mediterranean fattening operations already import 95 percent of the prey fish, mostly frozen sardines (Lovatelli 2003, Ottolenghi et al. 2004, Volpe 2005). In Mexico, more than half the catch of Pacific sardines is delivered directly to nearby tuna fattening pens, and much of the remaining catch is frozen or converted to fishmeal for use in other aquaculture operations. (Zertuche-González et al. 2008).



Steve Johnson

THREATS: CLIMATE CHANGE

Ocean predators are expected to suffer a wide range of impacts from global climate change, making them even more vulnerable to prey shortages. At the same time, climate change is also likely to affect prey species populations with changing temperatures, ocean currents, and sea ice.

Rising Temperatures

Prey fish and the food webs they support are highly sensitive to temperature changes, as seen in their dramatic population changes during El Niño and decade-long climate shifts (Anderson and Piatt 1999, Chavez et al. 2003). Small fish require favorable currents and temperatures to escape predators and to find enough to eat. The largest fishery in the world is vulnerable when warm El Niño waters bring Peruvian anchoveta toward the coast where they can be more easily caught. In 1972 this fishery crashed dramatically when heavy fishing coincided with an El Niño year (Clark 1976).

At the height of a 1950s cold period, temperature combined with heavy fishing to precipitate the collapse of the California sardine fishery made famous by John Steinbeck in his novel *Cannery Row*. Gulf of Alaska food webs are also tuned to temperature, and preferred prey such as capelin become scarce during warm periods, forcing marine mammals and seabirds to less nutritious options (Anderson and Piatt 1999). These prey shortages provide a preview of the likely effects of global warming.

Warm water in the North Sea brings fewer and smaller sand eels, the dominant prey for harbor porpoises (*Phocoena phocoena*), and could cause sand eels to hatch too soon for critical spring feeding time. Over the last thirty years, porpoises have suffered a 27 percent increase in starvation rates as sand eel populations shrink (Wanless et al. 2004, MacLeod et al. 2007a, 2007b). These climate-driven prey shortages are even more acute when prey populations are already overfished.



A puffin and an Arctic tern compete for prey

SEABIRDS AT RISK

Seabird populations are highly sensitive to their food supply, and increasingly threatened by human over-exploitation of prey fish. Crashes in fish populations in general have historically coincided with catastrophic seabird breeding failures, and climate-driven prey shortages may hit seabirds hardest.

When prey goes missing, seabirds rely on less nutritious alternatives and may become weak, delay their breeding season, and spend more time away from the nest searching for food (LeMaho et al. 1993). During herring declines in Norway's Lofoten Islands, the number of puffin burrows with chicks declined by 64 percent in the 1980s (Wright et al. 1996). Emaciated murrelets washed up on Norwegian shores after years of heavy fishing on capelin stocks, as populations of this usually common seabird plummeted by 56 to 80 percent (Hunt and Byrd 1999). In the eastern Bering Sea, both kittiwakes and murrelets starved to death during two decades of pollock shortages (Hunt and Byrd 1999, Springer et al. 1986).

50% The price increase in fish oil during El Niño in 1998.

— Tuominen and Esmark, 2003



Christopher Michel

Sea Ice

Polar predators and prey are under threat as the sea ice begins to melt (Smetacek and Nicol 2005, ACIA 2005). Krill aggregations feed on algae in a hidden sea ice ecosystem (Lizotte 2001, Loeb et al. 1997, Marschall 1988). As melting sea ice makes life harder in other ways, krill aggregations will shrink and in turn leave hungry penguins, whales, fish, and albatrosses in a weakened state. Warming and the decline of Antarctic krill have already been linked to reduced calving success of southern right whales that feed on this popular prey (Leaper et al. 2006)

Arctic predators are also threatened by the loss of sea ice and their prey, including narwhals, ringed seals, ivory gulls, and Atlantic salmon. Their favorite foods include Arctic cod (*Boreogadus saida*), which hides in ice cracks and channels and feeds on tiny organisms in the associated ecosystem (Bluhm and Gradinger 2008, Gradinger and Bluhm 2002, Tynan and DeMaster 1997).


Geographic Range Shifts

As ocean temperatures and currents are altered by climate change, the geographic ranges for predators and their prey are expected to move poleward. Wild Pacific salmon are predicted to shift northward, and sockeye salmon could shift entirely into the Bering Sea within the next 50 years (Welch et al. 1998, Mote et al. 2003). While some populations may adapt to their new location, others will likely suffer prey shortages. Salmon will be forced to seek out different prey, which may be less abundant in their new surroundings, leaving them malnourished and less able to adapt to other climate stresses (Bilby et al. 2007).



Wetjens Dimmlich

Despite their name, Crabeater seals primarily eat krill



Chris Dent

Penguins feeding krill to their chicks

SOLUTIONS: GETTING STARTED

Despite the alarming global outlook, some progress has been made toward the evaluation and management of prey species with predator needs in mind, as highlighted below. Successful management of predator-prey relations is an important first step toward ecosystem-based management, and must include prohibiting new fisheries for prey species, setting conservative catch limits for existing fisheries, prioritizing uses of prey species, and anticipating and reducing fishing during prey declines.

Manage for Ecosystem Integrity

Protecting prey is fundamental to keeping ocean ecosystems intact. State, regional, and federal agencies in the U.S. are beginning to recognize that healthy prey populations should be a goal for fishery managers. Yet initial efforts have devoted a disproportionate amount of time to gathering background information, with minimal actions toward preventing the actual loss of prey populations. Catch levels need to be set with large predatory fish, marine mammals, sea birds, and other marine life in mind.

The U.S. federal government is taking steps toward appropriate management of prey species. In 2002, federal regulations identified prey as a component of essential fish habitat and recognized the loss of prey as a problem for fisheries. More recently, regional fishery management councils have begun food web modeling with implicit consideration of predator needs.

States are also recognizing the importance of prey species. Most notably, Washington State was one of the first to create a dedicated management plan for forage fish and to explicitly adopt the protection of prey and “the integrity of the ecosystem and habitat upon which marine resources depend” as a goal for the Washington Department of Fish and Wildlife. Washington’s plan identifies knowledge gaps, addresses commercial and recreational catches, and describes the ecological role of prey fish. Along with Oregon and California, Washington has also banned fishing for krill, an important prey species. However, many states lag behind and in states with existing plans, management actions are urgently needed to actively protect prey fish.

Aziz T. Saltik



**Actively protect prey fish
rather than taking them for granted.**



Salvin's Albatross



Pablo Andrés Caceres Contreras

No New Fisheries for Prey Species

The most conservative approach to protecting prey is to ban fishing on these species before it begins. In 1998, the federal government banned fishing for prey species except pollock in federal waters around Alaska. The North Pacific Fishery Management Council and NOAA Fisheries acted to protect valuable groundfish and salmon predators targeted by fisheries, as well as other predators such as whales, sea lions, and albatrosses. The prey or “forage fish” protected from being caught included entire families of species such as krill (Euphausiacea), capelin and smelts (Osmeridae), sand lance (Ammodytidae) and many others.

Fishing for krill will likely similarly be banned in federal waters of the West Coast of the United States. The Pacific Fishery Management Council and NOAA Fisheries created a prohibited harvest species category to “ensure the long-term health and productivity of the West Coast ecosystem.” This proposed federal ban reinforces state bans on krill fishing established in California, Oregon, and Washington to prevent future exploitation of this important prey.

Set Conservative Catch Limits

Where fisheries on prey species already exist, conservative catch limits and careful monitoring are essential to prevent harm to natural predators. Squid, krill, and other prey fisheries sometimes remain unregulated or have incomplete management plans because they are not recognized as important.

Early management of Antarctic krill by the Convention on the Conservation of Antarctic Marine Living Resources (CCAMLR) exemplifies the conservative catch limits needed for prey species. With the ecosystem explicitly in mind, CCAMLR set initial catch limits for krill at nearly half the level of fishing mortality typically chosen for fisheries elsewhere in the world. Although there is substantial concern that this level is still too high, not fishing to the limit is an important start toward true ecosystem-based management.

Atlantic cod stocks have seen severe declines in recent decades and have yet to recover despite drastic cutbacks in fishing by the Canadian fleet (DFO 2005). Cod in the Northern Gulf of St. Lawrence also appear to be “starving to death” for unknown reasons which may include shortages of capelin, a preferred prey (DFO 2008, 2006). Conservative catch limits for capelin are an important component of Canada’s recovery strategy for cod populations in Newfoundland and Labrador (DFO 2006, 2005).



“To keep every cog and wheel is the first precaution of intelligent tinkering.”

— Aldo Leopold

Prioritize Uses for Prey

In addition to setting conservative catch levels, managers should give predators top priority before we divide up their prey, to better ensure that prey species fisheries are managed for the long-term health of the ecosystem. This includes specific assessment and allocation for the ecosystem - for fish, invertebrates, marine mammals, sea turtles, and other marine life.

After predator needs have been met, catches of prey species should be destined for direct human consumption. These small, fast-growing fish can reliably serve as primary protein for people. Peruvians have seen an upswing in anchoveta consumption since a 2006 event featuring top chef Gastón Acurio, scientist Patricia Majluf, and 18,000 diners feasting on flavorful fish that are more typically ground up for animal feed (Jacquet 2007).

Historically, direct consumption has been prioritized over use for bait, aquaculture, and other activities. California’s legislature in 1920 prohibited converting fish into oil and meal that would otherwise be fit for human consumption (Watson et al. 2006). The European Union later banned the use of Atlantic herring in fish meal in 2003, and England banned the use of fishmeal in ruminant feed due to concern over mad cow disease (Josupeit 2006, 2007). Peru banned the reduction of certain fish into oil and meal, including the South American pilchard (*Sardinops sagax*) and jurel or horse mackerel (*Trachurus murphyi*) (Watson et al. 2006).

Bait fish plays an important role in recreational and commercial fisheries, and should be prioritized after ecosystem and human needs. Aquaculture, animal feed, and all other uses for prey fish compete for these higher priority uses and require explicit acknowledgement and careful control. As aquaculture drives up prices for fishmeal and oil, prey fish now used for human consumption in developing regions are being diverted to industrial processing, with ethical implications (Alder and Pauly, 2006, FAO 2006, Delgado et al. 2003).

Fish Less During Natural Declines

Many prey species boom and bust with ocean temperature cycles, particularly sardines and anchovies. While it is normal for these populations to drop in response to changing conditions, heavy fishing during less productive phases could prevent the population from rebounding when favorable conditions return.

In the U.S., the Pacific Fishery Management Council is one of the first management organizations to explicitly account for the influence of ocean temperature cycles on sardines. This Council adopted a rule for managing Pacific sardine that adjusts the number of fish that can be caught according to sea surface temperature. Pacific sardines reproduce more slowly when the ocean is cooler, and fewer fish can be caught during this vulnerable time.

Sardines ready for human consumption





An Arctic tern catches its next meal

Paul McDonnell

SOLUTIONS: LOOKING AHEAD

While the needs of natural predators are increasingly recognized in public policy, concrete management actions to protect them are still generally lacking. However, several emerging science-based approaches could improve management of prey species substantially.

Maintain a Reserve for Natural Predators

Quantitative food requirements for each predator group are increasingly becoming available (Read and Brownstein 2003, Hunt and Furness 1996, Field and Francis 2006). Prey reserves can now be set aside for natural predators when catch limits are set for prey fish, squid, and krill.

At the stock assessment level, the needs of predators should be more rigorously accounted for in estimates of natural mortality and the definition of optimum yield for fisheries. At the policy and management level, explicit allocations can be made for ecosystem needs or for natural predators.



Wetjens Dimmlich

Female elephant seal and pup

Protect Breeding Hotspots

For predators with clear needs for prey during spawning, nursing, or nesting season, closures on fishing for squid, anchovy, or other prey may be appropriate. This would prohibit fishing in very specific times and places to increase prey availability and reproductive success with the added benefit of reducing the number of predators caught in the nets. Unfortunately the specific needs of many predators remain unknown.

In Punta Tombo, Argentina, tracking experiments with Magellanic penguins clearly show their most important foraging areas in need of protection (Boersma 2008). In the North Sea, fishing activity and seabird feeding are concentrated in different locations suggesting the potential for successful zoning (Furness and Tasker 2000, Hunt and Furness 1996). Separating fishing from seabird feeding could minimize fishery impacts and prevent localized depletion of prey.

Save 10% for Climate Losses

In addition to planning for ecosystem needs, managers should leave a buffer for climate-driven losses in prey populations. In the absence of more quantitative decision rules based on temperature, managers should insure against climate change impacts with a reduction of ten percent in catch limits.

Manage in Real Time

Real-time monitoring during the fishing season is especially helpful for management of short-lived and infrequently counted prey fish and invertebrates. Daily updates on catch and effort for each vessel allow managers to monitor the status of the fishery and decide whether too many fish have been taken in a season. Real-time monitoring is currently active for the squid (*Loligo gahi*) fishery in the Falkland Islands and has been proposed for United States squid fisheries (Agnew et al. 1998). Real-time monitoring of fishing intensity has also been proposed to protect Peruvian anchoveta, which are more vulnerable to being caught as they move closer to shore during El Niño climate cycles (Bertrand et al. 2005).

SOLUTIONS SUMMARY

GOAL:

Protect prey species as fundamental to ocean ecosystems and fisheries.

SOLUTIONS:


1. No new fisheries for prey
2. Set conservative limits for prey fisheries and save 10 percent for climate impacts
3. Protect breeding hotspots
4. Prioritize uses for prey—place ecosystem needs first






PREY SPECIES AROUND THE WORLD


Anchovy




Squid



Krill




Prey Fish





More Prey Fish



Dolphins and bait ball

RESOURCES

Abitia-Cardenas LA, Galvan-Magana F, Gutierrez-Sanchez J, Rodriguez-Romero J, Aguilar-Palomino B, Moehl-Hitz A. (1999) Diet of blue marlin *Makaira mazara* off the coast of Cabo San Lucas, Baja California Sur, Mexico. *Else*, 44, 95-100

ACIA. 2005. Arctic Climate Impact Assessment. Cambridge University Press.

Alder J, B. Campbell, V. Karpouzi, K. Kaschner, and D. Pauly. 2008. Forage fish: From ecosystems to markets. *Annual Reviews in Environmental Resources* 33:7.1-7.14.

Alder J, Pauly D (2006) Human consumption of forage fish. In: Alder J, Pauly D (EDs.) *On the multiple uses of forage fish: from ecosystems to markets*. Fisheries Centre Research Reports 14(3) Fisheries Centre, University of British Columbia

American Cetacean Society. Fact Sheet: Blue Whale. Accessed 3 October 2007. <http://www.acsonline.org/factpack/bluewhl.htm>

American Heart Association (AHA). 2007. Fish and Omega-3 Fatty Acids. <http://www.americanheart.org/presenter.jhtml?identifier=4632>

Associated Press (2001) Giant Tuna Sells for Record \$173,600. <http://www.flmnh.ufl.edu/fish/InNews/GiantTuna.htm>

Arzel P. et al., Les algues et invertébrés marins des pêches françaises—2eme partie, Algues, Eponges, Corail rouge, Escargots de mer, Bivalves, Poulpes, Seiches, Encornets,

Atkinson A, Siege V, Pakhomov E, Rothery P. (4 November 2004) "Long-term decline in krill stock and increase in salps within the Southern Ocean." *Nature* 432, 100-103.

The Atlantic Menhaden Plan Review Team (2004), 2004 Review of the Fishery Management Plan for Atlantic Menhaden (*Brevoortia Tryannus*).

Atlantic States Marine Fisheries Commission (2003) Species Profile: Atlantic Herring - Atlantic Herring Fishery Management Plan <http://www.asmfc.org/speciesDocuments/herring/herringProfile.pdf>

Australian Government Antarctic Division, Krill <http://www.aad.gov.au/default.asp?casid=1540> [Anonymous] Australian Marine Conservation Society, Farming Australia's Seas; It's No Solution! http://www.amcs.org.au/default2.asp?active_page_id=159

Baranowski R, Beddington J.R., des Clers S, Nolan CP (1998) Approaches to assessing stocks of *Loligo gahi* around the Falkland Islands. *Fisheries Research* 35: 155-169

Bartlett L. (1994) Alaska Department of Fish and Game; Eulachon. <http://www.adfg.state.ak.us/pubs/notebook/fish/eulachon.php>

Bargmann G (1998) Forage Fish Management Plan. Washington Fish and Wildlife Commission, 24 January 1998. <http://wdfw.wa.gov/fish/forage/manage/foragman.pdf>

Bearzi, G., R.R. Reeves, G. Notarbartolo-Di-Sciara, E. Politi, A. Cañadas, A. Frantzis, B. Mussi. (2003) Ecology, status and conservation of short-beaked common dolphins, *Delphinus delphis*, in the Mediterranean Sea. *Mammal Review* 33(3-4):224-252.

Becker, B.H. and S.R. Beissinger. 2006. Centennial decline in the trophic level of an endangered seabird after fisheries decline. *Conservation Biology* 20(2):470-479.

Bell, Tom. (2006). Offshore Opportunity? *Portland Press Herald*. December 17, 2006. <http://www.choircoalition.org/media/detail.php?id=24>

Bertrand S, Burgos JM, Gerlotto F, Atiquipa J (2005) Lévy trajectories of Peruvian purse-seiners as an indicator of the spatial distribution of anchovy (*Engraulis ringens*). *ICES Journal of Marine Science* 62: 477-482

Bilby R., Hannah S., Healey M., Huntley N., Hurlbert S., Lamberson R., Levings C., Montgomery D., Percy W., Poe T. P., Smouse P. "Climate Change Impacts on Columbia River Basin Fish and Wildlife." Independent Scientific Advisory Board. p.90.

Block BA, Teo SLH, Walli A, Boustany A, Stokesbury MJW, Farwell CJ, Weng KC, Dewar H, Williams TD (2005) Electronic tagging and population structure of Atlantic bluefin tuna. *Nature* 434: 1121-1127

Blue Ocean Institute. Species score cards: Argentine squid <http://www.blueocean.org/seafood/species/21.html>

Blue Ocean Institute - Species score cards: Longfin squid <http://www.blueoceaninstitute.org/seafood/species/126.html>

Blue Ocean Institute - Species score cards: Market squid <http://www.blueocean.org/seafood/species/122.html>

Blue Ocean Institute - Species score cards: Shortfin squid <http://www.blueocean.org/>

[seafood/species/103.html](http://www.blueocean.org/seafood/species/103.html)

Bлум, B.A. and R. Gradinger. (2008) Regional Variability In Food Availability For Arctic Marine Mammals. *Ecological Applications* 18(2) Supplement: S77-S96.

Bochenek E.A.. *Ilex and Loligo Squid Real-Time Management Studies*. Haskin Shellfish Research Laboratory, Rutgers University. <http://www.hsrl.rutgers.edu/squid.html>

Boersma, D. (2008) Penguins as marine sentinels. *BioScience* 58(7):597-607.

Brame D, Duval M, Goldsborough, Hinman, Ken and Schick, Amy Menhaden Matter, Chesapeake Bay Foundation, Coastal Conservation Association, Environmental Defense, and National Coalition for Marine Conservation.

Brander K (2006) "Assessment of possible impacts of climate change on fisheries." German Advisory Council on Climate Change, ICES Copenhagen. http://www.wbgu.de/wbgu_sn2006_ex02.pdf

Brodeur RD, Daly EA, Schabetsberger RA, Mier KL (2007) Interannual and interdecadal variability in juvenile coho salmon (*Oncorhynchus kisutch*) diets in relation to environmental changes in the northern California Current. *Fisheries Oceanography* 16(5): 395-408

Bronsgest-Schoute HC, van Gent CM, Luten JB, Ruiters A. The effect of various intakes of omega 3 fatty acids on the blood lipid composition in healthy human subjects. *Am J Clin Nutr*. 1981 Sep; 34 (9):1752-1757.

Bruggeman P. Underwater Field Guide, Ice krill *Euphausia crystallorophias* <http://scilib.ucsd.edu/sio/nsf/fguide/arthropoda10.html>

Buschmann, A.H., M.C. Hernández González, L. Henríquez Antipa. 2006. *Salmonicultura y su dependencia de la pesca en Chile*. Documento 22, Oceana.

Butler CM (2007) Atlantic Bluefin Tuna (*Thunnus thynnus*) Feeding Ecology and Potential Ecosystem Effects During Winter in North Carolina. Master's Thesis: North Carolina State University.

Campbell, B., Alder, J. (2006) "Fishmeal and fish oil: production trade and consumption," In: Alder, J., Pauly D. (eds). "On the multiple uses of forage fish: From ecosystems to markets." Fisheries Centre Research Reports 14(3). Fisheries Centre, University of British Columbia [ISSN 1198-6727]



- Chavez F P., John Ryan, Salvador E. Lluch-Cota and Miguel Niguen C. (2003) From anchovies to sardines and back: Multidecadal change in the Pacific Ocean. *Science* 299: 217-221.
- Chase BC (2002) Differences in diet of Atlantic bluefin tuna (*Thunnus thynnus*) at five seasonal feeding grounds on the New England continental shelf. *Fisheries Bulletin* 100: 168-180
- Clapham P.J., Young SB, Brownell Jr. RL. (1999) Baleen whales: conservation issues and the status of the most endangered populations. *Mammal Review* 29, 35-60
- Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR). CCAMLR Ecosystem Approach <http://www.ccamlr.org/pu/e/sc/eco-app-intro.htm>
- Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR). CCAMLR's Approach to Management http://www.ccamlr.org/pu/e/e_pubs/am/toc.htm
- Commission for the Conservation of Antarctic Marine Living Resources (CCAMLR). CCAMLR: Harvested Species: Krill (*Eupausia superba*) <http://www.ccamlr.org/pu/E/sc/fish-monit/hs-krill.htm>
- Croxall J.P., Trathan P.N., Murphy E.J. (30 August 2002) Environmental Change and Antarctic Seabird Populations. *Science* 297: 1510-1514
- Cury, P., A. Bakun, R.J.M. Crawford, A. Jarre-Teichmann, R.A. Quiñones, L.J. Shannon, and H.M. Verheye. 2000. Small pelagics in upwelling systems: patterns of interaction and structural changes in "wasp-waist" ecosystems. *ICES Journal of Marine Science* 57:603-613.
- Defenders of Wildlife. Navigating the Arctic Meltdown: Arctic Cod. http://www.defenders.org/resources/publications/programs_and_policy/science_and_economics/global_warming/navigating_the_arctic_meltdown_arctic_cod.pdf Accessed 15 October 2007.
- Delgado, C.L., N. Wada, M.W. Rosegrant, S. Meijer, and M. Ahmed. (2003) *Fish to 2020: Supply and Demand in Changing Global Markets*. International Food Policy Research Institute and The World Fish Center.
- Department of Fisheries and Oceans, Canada (DFO). (2006) Press Release "Minister Hearn announces 2006 management plan for 2J3KL, 3Ps and 4RST capelin fisheries." http://www.nfl.dfo-mpo.gc.ca/publications/news_presse_NL_TNL_0622.asp
- Department of Fisheries and Oceans, Canada (DFO). (2008) What's Holding Back the Cod Recovery? http://www.dfo-mpo.gc.ca/science/Story/quebec/cod_recovery_e.htm
- Department of Fisheries and Oceans, Canada (DFO). (2005) Canada-Newfoundland and Labrador Action Team for Cod Recovery. A Strategy for the Recovery and Management of Cod Stocks in Newfoundland and Labrador. http://www.dfo-mpo.gc.ca/cod-morue/Strategie_nl_e.htm
- Dery, L.M. (2004) Atlantic herring, *Clupea harengus*, NOAA's National Marine Fisheries Service, Northeast Fisheries Science Center <http://www.nefsc.noaa.gov/fbi/age-man/her/her.htm>
- Environmental Defense (2005) Why Do Atlantic Menhaden Matter? <http://www.environmentaldefense.org/article.cfm?contentID=4561>
- Estes JA, T. T., Doak DF (1998) Killer Whale Predation on Sea Otters Linking Oceanic and Nearshore Ecosystems. *Science* 282: 473-476.
- Eurofish (2005) "FISH INFOnetwork Market Report—Surimi." <http://www.eurofish.dk/indexSub.php?id=1928&easysitestatid=-1765644808>
- FAO (2004) Species Fact Sheet, *Engraulis ringens*. <http://www.fao.org/figis/servelet/species?fid=2917>
- FAO (2004) The State of World Fisheries and Aquaculture, 2004 Part 1: World review of fisheries and aquaculture, The status of fishery resources. http://www.fao.org/DOCREP/007/y5600e/y5600e05.htm#P1235_55689
- FAO (2005) Fisheries Report No. 779 (FIRI/R779) Report of the Third Meeting of the AD HOC GFCM/ICCAT Working Group on Sustainable Bluefin Tuna Farming / Fattening Practices in the Mediterranean, General Fisheries Commission for the Mediterranean international Commission of the Conservation of Atlantic Tunas, Rome, 16-18 March 2005 <http://www.fao.org/docrep/008/y8870e/y8870e00.htm>
- FAO (2005) Review of the state of world marine fishery resources. <ftp://ftp.fao.org/docrep/fao/007/y5852e/Y5852E23.pdf>
- FAO (2006) State of the world fisheries and aquaculture: Trash or Treasure? Low-value/trash fish from marine fisheries in Asia-Pacific region. Food and Agriculture Organization of the United Nations (FAO) <http://www.fao.org/docrep/009/A0699e/A0699E07.htm>
- FAO. Animal Feed Resources Information System, Animal Products <http://www.fao.org/ag/aga/AGAP/FRG/afris/DATA/332.htm>
- FAO. Fish Synop., (125)Vo1.7,Pt.2:305-579 FAO Species catalogue Vol. 7. Clupeoid fishes of the world. (Suborder CLUPEOIDEI)
- FAO Marine Resources Service, Fishery Resources Division. Review of the state of world marine fishery resources FAO Fisheries Technical Paper. No.457. Rome, FAO. 2005. 235 <ftp://ftp.fao.org/docrep/fao/007/y5852e/Y5852E00.pdf>
- FAO Species Fact Sheet <http://www.fao.org/fi/website/FIRetrieveAction.do?dom=species&fid=3017>
- FAO, Species Fact Sheet: *Loligo pealei*. <http://www.fao.org/fi/website/FIRetrieveAction.do?dom=species&fid=2714>
- FAO/SIDP Species Identification Sheets: *Illex illecebrosus* http://www.oceansatlas.org/world_fisheries_and_aquaculture/html/resources/capture/mainspec/species/sp2720.htm
- FAO/SIDP Species Identification Sheets: *Illex argentinus* http://www.oceansatlas.org/world_fisheries_and_aquaculture/html/resources/capture/mainspec/species/sp3565.htm
- FAS Online, (2000) Special Report- December 2000, BSE in Europe: The Impact on U.S. and World Soybean and Soybean Meal Demand <http://www.fas.usda.gov/oilseeds/circular/2000/00-12/decspec.htm>
- Federal Register (2007) <http://edocket.access.gpo.gov/2007/pdf/e7-3247.pdf>
- Field JC and RC Francis. (2006) Considering ecosystem-based fisheries management in the California Current. *Marine Policy* 30:552-569.
- FIGIS—Fisheries Statistic Production (FAO) <http://www.fao.org/fi/website/FIRetrieveAction.do?dom=collection&xml=global-production.xml>
- FIRMS Marine Resource Fact sheet: Squid - Global, 2005 <http://www.fao.org/figis/website/FIRMSRetrieveAction.do?dom=resource&fid=13335>
- FishBase. Family *Engraulidae*—Anchovies <http://www.fishbase.org/Summary/FamilySummary.cfm?Family=Engraulidae>

- FishBase. Alaska Pollock.
<http://www.fishbase.org/Summary/SpeciesSummary.php?id=318>
- Fishmeal Information Network (FIN), Fishmeal Facts and Figures <http://www.gafta.com/fin/finfacts1.html>
- Fishmeal Information Network (FIN) Fishmeal and Fish Oil in a safe and sustainable food chain (2002) <http://www.gafta.com/fin/MEPBriefing.pdf>
- Forcada J, Trathan PN, Reid K, Murphy EJ. (2005) "The Effects of Global Climate Variability in Pup Production of Antarctic Fur Seals." *Ecology* 86(9)
- Forcada J, Trathan PN, Reid K, Murphy EJ, Croxall JP. (2006) Contrasting population changes in sympatric penguin species in association with climate warming. *Global Change Biology* 12: 411-423
- Franklin B. (2001). The Most Important Fish in the Sea. *Discover*. September 2001.
- Frederiksen M, Wanless S, Harris MP, Rothery P, Wilson LJ. (2004) The role of industrial fisheries and oceanographic change in the decline of North Sea black-legged kittiwakes. *Journal of Applied Ecology* (41) 1129-1139.
- Furness, R.W. and M.L. Tasker. (2000) Seabird-fishery interactions: quantifying the sensitivity of seabirds to reductions in sandeel abundance and identification of key areas for sensitive seabirds in the North Sea. *Marine Ecology Progress Series* 202:253-264.
- Golet WJ, Cooper AB, Campbell R, Lutcavage M (2007) Decline in condition of northern bluefin tuna (*Thunnus thynnus*) in the Gulf of Maine. *Fisheries Bulletin* 105(3): 390-395
- Gómez-Gutiérrez, J., Euphausiids <http://www.geocities.com/jgomez64/euphausiids.html>
- Government of Newfoundland and Labrador (2005) A Strategy for the Recovery and Management of Cod Stocks in Newfoundland and Labrador.
http://www.dfo-mpo.gc.ca/cod-morue/nl_e.pdf
- Grabowski, Jonathan H. (2003) Are We Using Herring to Farm Lobsters? The Effects of Herring Bait on Diet Composition and Growth of the American Lobster The Lobster Bulletin' <http://kodiak.asap.um.maine.edu/lobster/library/publications/bulletin/summer2003.htm#herring>
- Gradinger R. and B. Bluhm. (2002) Life in cold gaps. NOAA Ocean Explorer Explorations. Arctic Exploration. <http://oceanexplorer.noaa.gov/explorations/02arctic/logs/sep3/sep3.html>
- Griffin JC, Magraf FJ (2003) The diet of Chesapeake Bay striped bass in the late 1950s. *Fisheries Management and Ecology* 10: 323-328
- Guenneugues, P., Morrissey, M. 2005. "Surimi Resources." In: Park, J.W. (Ed). *Surimi and Surimi Seafood*, second edition. Marcel Dekker, Inc. Basil Switzerland, 2005.
- Hartman KJ, Magraf FJ (2003) US Atlantic coast striped bass: issues with a recovered population. *Fisheries Management and Ecology* 10: 309-312
- Heckle, Harold (2006) *Mass Beaching Piles on the Anchovies*, Associated Press Available from: http://www.livescience.com/animalworld/060929_ap_anchovie_beaching.html
- Herbst, S. 1995. "Surimi." *International recipes: Food and Wine Dictionary*. Available from <https://www.internationalrecipesonline.com/recipes/dictionary.pl?6729>
- Helga Josupeit (2006) FISH INFO Network Market Report, FAO
- Hendrickson, L. Northern shortfin squid (*Illex illecebrosus*)
 Status of Fishery Resources off the Northeastern US - NEFSC - Resource Evaluation and Assessment Division, NOAA
<http://www.nefsc.noaa.gov/sos/spsyn/iv/sfsquid/>
- Howard, D. Krill in Cordell Bank National Marine Sanctuary, NOAA. <http://oceanexplorer.noaa.gov/explorations/02quest/background/krill/krill.html>
- Hucke-Gaete R, Viddi F, Bell M (2006) *Marine Conservation in southern Chile*. Impreso en Imprenta America, Valdivia, Chile. 124 pp.
- Hucke-Gaete R., Osman L.P., Moreno C.A., Findlay K.P., Ljungblad D.K. (2003) Discovery of blue whale feeding and nursing ground in southern Chile. *Proceeding of the Royal Society of London B (Suppl.)*, Biology Letters.
- Hunt G.L. and Byrd G.V. (1999) Marine Bird Populations and Carrying Capacity of the Eastern Bering Sea. *Dynamics of the Bering Sea*. (631-650)
- Hunt, GL, Jr. and RW Furness (eds.) (1996) *Seabird/fish interactions, with particular reference to seabirds in the North Sea*. ICES Cooperative Research Report No. 216. International Council for the Exploration of the Sea, Copenhagen.
- Hunt SL, Mulligan TJ, Komori K (1999) Oceanic feeding habits of chinook salmon *Oncorhynchus tshawytscha*, off northern California. *Fishery Bulletin* 97(3): 717-721
- ICCAT. (2006) Atlantic Bluefin Tuna Stock Assessment.
<http://www.iccat.int/assess.htm>
- IFREMER. (1992) *Crustacés, Oursins et Violets*, IFREMER, 135p. <http://www.ifremer.fr/docelec/doc/1992/rapport-1535.pdf>
- Industry Canada (1996) The Role of Capelin, <http://www.cdli.ca/cod/role.htm>
- Information Center for the Icelandic Ministry of Fisheries <http://www.fisheries.is/stocks/capelin.htm>
- International Council for the Exploration of the Sea (ICES). (2004) Blue Whiting: playing a big game with small fish. *ICES Newsletter*. September 2004.
- International Fishmeal and Fish Oil Organization (IFFO), Datasheet: The production of fishmeal and fish oil from gulf menhaden.
- IPPC (2007) Summary for Policymakers. In: *Climate Change 2007: Climate Change Impacts, Adaptation and Vulnerability*. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.
- Jackson, J.B.C. 2008. Ecological extinction and evolution in the brave new ocean. *Proceedings of the National Academy of Sciences* 105 supplement 1: 11458-11465.
- Jacobson, L.D. (2005) *Essential Fish Habitat Source Document: Longfin Inshore Squid, Loligo pealeii, Life History and Habitat Characteristics (Second Edition)*, NOAA Technical Memorandum NMFS-NE-193, September 2005.
<http://www.nefsc.noaa.gov/nefsc/publications/tm/tm193/tm193.pdf>
- Jacquet J (2007) Save Our Oceans, Eat Like a Pig. The Yee 17 April 2007. <http://theyee.ca/Views/2007/04/17/EatLikePigs/> Accessed: 14 September 2007
- Jennings S, Kaiser MJ, Reynolds JD (2001) *Marine Fisheries Ecology*. Blackwell Publishing: Malden, MA. 417p.
- Josupeit, H. (2006). Fish oil Market Report—March 2006. FAO GlobeFish. <http://www.globefish.org/dynamisk.php4?id=3654>



Josuweit, H. (2007). Fishmeal Market Report—April 2007. FAO GlobeFish. <http://www.globefish.org/dynamisk.php?id=3654>

Josuweit, H. 2006. "Fishmeal Market Report—June 2006." FAO GlobeFish. <http://www.globefish.org/index.php?id=3015>

Josuweit, H. 2007. "Fishmeal Market Report—September 2007." FAO GlobeFish. <http://www.globefish.org/index.php?id=4286>

June, F and J.W. Reintjes, J. The menhaden fishery. In: M.E. Stansby, Editor, Industrial Fishery Technology, R.E. Krieger Publishing Company, NY (1976), pp. 136–149.

Kaschner K, P. D. (2004). Competition between marine mammals and fisheries: food for thought. T. H. S. International. Vancouver, Canada, MMRU Fisheries Centre: 1-28.

Kassahn KS, Caley MJ, Ward AC, Connolly AR, Stone G, Crozier RH. (2007) "Heterologous microarray experiments used to identify the early gene response to heat stress in a coral reef fish." *Molecular Ecology* 16, 1749-1763.

Kaschner, K., Karpouzi, V., Watson, R. Pauly, D. 2006. "Forage Fish consumption by marine mammals and seabirds." In: Alder, J., Pauly D. (eds). "On the multiple uses of forage fish: From ecosystems to markets." Fisheries Centre Research Reports 14(3). Fisheries Centre, University of British Columbia [ISSN 1198-6727]

Kelly, K.H., and J.R. Moring. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North Atlantic)—Atlantic herring. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.38). U.S. Army Corps of Engineers, TR EL-82-4. 22 pp.

Koslow J. A., G. W. B., J. D. M. Gordon, and P. L. R. L. Haedrich, and N. Parin (2000). "Continental slope and deep-sea fisheries: implications for a fragile ecosystem." *ICES Journal of Marine Science* 57: 548-557.

Kucas, S.T., Jr. 1986. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (Pacific Southwest)—northern anchovy. U.S. Fish Wildl. Serv. Biol. Rep. 82(11.50). U.S. Army Corps of Engineers, TR EL-82-4. <http://www.nwrc.usgs.gov/wdb/pub/0142.pdf>

Leaper R, Cooke J, Trathan P, Reid K, Rowntree V, Payne R. (2006) "Global climate drives southern right whale (*Eubalaena australis*) population dynamics." *Biol. Lett.* 2, 289–292.

Learmonth, J.A., C.D. Macleod, M.B. Santos, G.J. Pierce, H.Q.P.Crick, and R.A. Robinson. (2006)

Potential Effects of Climate Change on Marine Mammals. *Oceanography and Marine Biology: An Annual Review* 44:431-464

Lewis A.D. and P Dalzell. (1989). A Review of the South Pacific Tuna Baitfisheries: small pelagic fisheries associated with coral reefs. *Marine Fisheries Review*. 51 n4

LeMaho Y., Gendner J.P., Challet E., Bost C.A., Gilles J., Verdon C., Plumeré C., Robin J.P., Handrich Y(1993). Undisturbed breeding penguins as indicators of changes in marine resources. *Marine Ecology Progress Series* 95: 1-6

Lilli Ann, LLC <http://sec.edgar-online.com/2004/07/20/0001193125-04-120537/Section19.asp>

Loeb, V., V. Siegal, O. Holm-Hansen, R. Hewitt, W. Fraser, W. Trivelpiece, and S. Trivelpiece. (1997) Effects of sea-ice extent and krill or salp dominance on the Antarctic food web. *Nature* 387: 987-900.

Lovatelli (2003) Summary Report on the Status of BFT Aquaculture. In Report of the 2nd Meeting of the ad hoc Working Group on Sustainable Tuna Farming/Fattening Practices in the Mediterranean, Izmir, Turkey, 15-17 December 2003, pp. 73-89. GFC and ICCAT <http://www.fao.org/docrep/008/y8870e/y8870e0e.htm#bm14>

Luna, Susan M. and Alexi Orlov (2005) "Mallotus villosus", Capelin, FishBase 10 2005 Version

MacLeod C, Santos M, Reid RJ, Scott B, Pierce GJ. (2007a) "Linking sand eel consumption and the likelihood of starvation in harbour porpoises in the Scottish North Sea: could climate change mean more starving porpoises?" *Biol. Lett.* 3, 185–188.

MacLeod CD, Pierce GJ, Santos MB. (2007b) "Starvation and sandeel consumption in harbour porpoises in the Scottish North Sea." *Biol. Lett.* 3, 535–536.

Mahmoudi, B., McBride, R. 2002. "A review of Florida's halfbeak bait fishery and halfbeak biology and a preliminary stock assessment." Florida Fish and Wildlife Conservation Commission.

Manooch CS III (1973) Food Habits of Yearling and Adult Striped Bass, *Morone saxatilis* (Walbaum), from Albemarle Sound, North Carolina. *Chesapeake Science* 14(2): 73-86

Marine Life in Whatcom County, Fish Species: Pacific Sand Lance (*Ammodytes hexapterus*) http://whatcommrc.wsu.edu/Fact_Sheets/sandlance.htm

Marine Institute of Memorial University of Newfoundland—Offshore/inshore Fisheries Development. <http://www.mi.mun.ca/mi-net/fishdeve/capelin.htm>

Market Report 2004-2005. http://www.st.nmfs.gov/st1/market_news/doc76.txt

Marschall, H-P. (1988) The overwintering strategy of Antarctic krill under the pack-ice of the Weddell Sea. *Polar Biology* 9(2):129-135.

Menhaden Resource Council. <http://www.menhaden.org>

Montague, P (2003) Australian southern bluefin tuna farming and research activity—national report. *Cahiers Options Méditerranéennes* 60:139-141

Monterey Bay Aquarium Seafood Watch http://www.mbayaq.org/cr/SeafoodWatch/web/sfw_factsheet.aspx?fid=22

Morato T., W. R., Pitcher TJ, Pauly D. (2006). "Fishing down the deep." *Fish and Fisheries* 7: 24-34.

Morgan J, Hiller M. "Are we putting our fish in hot water?" World Wildlife Fund Climate Change Programme. http://www.panda.org/about_wwf/what_we_do/climate_change/index.cfm?uNewsID=50460

Mote PW, Parson EA, Hamlet AF, Keeton WS, Lettenmaier D, Mantua N, Miles EL, Peterson DW, Peterson DL, Slaughter R, Snover AK. (2003) "Preparing for Climatic Change: The Water, Salmon, and Forests of the Pacific Northwest." *Climatic Change* 61, 45-88.

Naylor, R.L., R.J. Goldburg, J.H. Primavera, N. Kautsky, M.C.M. Beveridge, J. Clay, C. Folke, J. Luchenco, H. Mooney, and M. Troell. (2000) Effect of Aquaculture on World Fish Supplies. *Nature*. Vol. 405

New, M.B. and U.N. Wijkstrom. (2002). Use of fishmeal and fish oil in aquafeeds: Further thoughts on the fishmeal trap. Food and Agriculture Organization of the United Nations (FAO)

Nicol, Stephen and Yoshinari Endo (1997) Krill Fisheries of the World, Fisheries Technical Paper 367, FAO <http://www.fao.org/docrep/003/w5911e/w5911e00.htm>

Nicol, Stephen and Jacqueline Foster (2003) Recent Trends in the Fishery for Antarctic Krill, *Aquatic Living Resources*. 16:42–45

NOAA. Collapse of Anchovy Fisheries and the Expansion of Sardines In Upwelling Regions. <http://www.pfeg.noaa.gov/research/climatemarine/cmffish/cmffishery4.html>

Norway's official site on seafood safety and resource and aquaculture management, Marine Stocks, North-East Arctic Cod. http://www.fisheries.no/marine_stocks/fish_stocks/cod/north_east_arctic_cod.htm

Office of Science and Technology (2006) Production of Fish Meal and Fish Oil, NOAA Fisheries http://www.st.nmfs.gov/st1/market_news/doc76.txt

Office of Science and Technology (2006) Foreign Trade in Industrial Products, NOAA Fisheries http://www.st.nmfs.gov/st1/market_news/doc76b.txt

Ottolenghi, F., C. Silvestri, P. Giordano, A. Lovatelli, M.B. New (2004) Capture-Based Aquaculture: The Fattening of eels, groupers, tunas, and yellowtails. FAO.

Pacific Seafood Group http://www.pacseafood.com/products/al_pollock.html

Pacific States Marine Fisheries Commission. http://www.psmfc.org/habitat/edu_anchovy_fact.html

Paine RT, Ruesink JL, Sun A, Soulanille EL, Wonham MJ, Harley CDG, Brumbaugh DR, Secord DL (1996) Trouble on Oiled Waters: Lessons from the Exxon Valdez Oil Spill. *Annual Review of Ecology and Systematics* 27: 197-235

Pauly D., C. V., Dalsgaard J., Froese R., Torres F. (1998) Fishing down marine food webs. *Science* 279: 860-864.

Pauly, D. (1992) Peruvian anchoveta, Charles Darwin and us. *Naga, the ICLARM* 15(4):14-15. <http://data.fisheries.ubc.ca/references/Darwin/199217-PeruAnchoveta.pdf>

Perez Matus AA, Buschmann AH (2003) Sustentabilidad e Incertidumbre de las Principales Pesquerías Chilenas. *Oceana*: Santiago, Chile. 114p.

Peterson CH, Rice SD, Short JW, Esler D, Bodkin JL, Ballachey BE, Irons DB (2003) *Science* 302: 2082-2086

PFMC Amendment 12 <http://www.pcouncil.org/cps/cpsfmp/cpsa12.html>

Piatt, J. and W. Sydeman. (Unpublished) Seabirds as Indicators of Marine Ecosystems: A Brief History.

Politi E., B. G., Airoldi S. (2000). "Evidence for malnutrition in bottlenose dolphins photoidentified in the eastern Ionian Sea." *European Research on Cetaceans* 14: 234-236.

Powell, Kendall (2003) *Eat Your Veg.* Nature Vol. 426

Read, A. J. and C. R. Brownstein. 2003. Considering other consumers: fisheries, predators, and Atlantic herring in the Gulf of Maine. *Conservation Ecology* 7(1): 2. <http://www.consecol.org/vol7/iss1/art2/>

Remarks of William J. Goldsborough; Chesapeake Bay Foundation Senior Scientist (May 26, 2004) to the Atlantic Menhaden Management Board of the Atlantic States Marine Fisheries Commission.

Rodhouse, P.G.; Dawe, E.G.; O'Dor, R.K. (EDs) Squid recruitment dynamics. The genus *Illex* as a model. The commercial *Illex* species. Influences on variability. FAO Fisheries Technical Paper. No. 376. Rome, FAO. 1998. 273p. <ftp://ftp.fao.org/docrep/fao/008/w9000e/w9000e00.pdf>

Roessig JM, Woodley CM, Cech Jr. JJ, Hansen LJ. (2004) "Effects of global climate change on marine and estuarine fishes and fisheries." *Reviews in Fish Biology and Fisheries* 14: 251-275.

Roper, C.F.E., M.J. Sweeney, C.E. Nauen. 1984. FAO Species catalogue: Cephalopods of the world An Annotated and Illustrated Catalogue of Species of Interest to Fisheries. FAO Fisheries Synopsis No. 125, Volume 3

Rosen DAS, Trites AW (2000) Pollock and the decline of Stellar sea lions: testing the junk-food hypothesis. *Canadian Journal of Zoology* 78: 1243-1250

Royal Society for the Protection of Birds (RSPB) Demand for fish meal and fish oil <http://www.rspb.org.uk/policy/marine/fisheries/sustainable/demand.asp>

Ruttan, L. and Tyedmers, P. 2007. Skippers, spotters and seiners: Analysis of the "skipper effect" in US menhaden (*Brevortia* spp.) purse-seine fisheries. *Fisheries Research* 83:73-80

Sandvik H, Erikstad KE, Barrett RT, Yoccoz NG. (2005) "The effect of climate on adult survival in five species of North Atlantic seabirds." *Journal of Animal Ecology* 74: 817-831

Sarà, G. and R. Sarà (2007) Feeding habits and trophic levels of bluefin tuna *Thunnus thynnus* of different size classes in the Mediterranean Sea. *Journal of Applied Ichthyology* 23(2):122-127

Shepherd, C.J., I.H. Pike, and S.M Barlow. (2005) Sustainable Feed Resources of Marine Origin. *European Aquaculture Society Special Publication* June 2005. 35: 59-66

Sinopoli, M., C. Pipitone, S. Campagnuolo, D.Campo, L.Castriota, E. Mostarda, F. Andaloro. 2004. Diet of young-of-the-year bluefin tuna, *Thunnus thynnus* (Linnaeus, 1758), in the southern Tyrrhenian (Mediterranean) Sea. *Journal of Applied Ichthyology* 20(4):310-313

Skewgar, Elizabeth; Boersma, P. Dee; Harris, Graham; Caille, Guillermo (2007) Anchovy Fishery Threat to Patagonian Ecosystem, *Science*, Vol. 315

Smetacek V, Nicol S. (15 September 2005) Polar ocean ecosystems in a changing world. *Nature* 437: 362-368

Species Fact Sheet: *Engraulis mordax* <http://www.fao.org/figis/servlet/species?fid=2107>

Species Fact Sheet: *Engraulis anchoita* <http://www.fao.org/figis/servlet/species?fid=2916>

Springer A.M., Roseneau D.G., Lloyd D.S., McRoy C.P., Murphy E.C. (1986). Seabird responses to fluctuating prey availability in the eastern Bering Sea. *Marine Ecology Progress Series* 32: 1-12

Stephanie Showalter, The National Sea Grant Law Center, American Pelagic Fishing Company v. U.S., 55 Fed. Cl. 575 (2003). <http://www.olemiss.edu/orgs/SGLC/National/SandBar/2.2taking.htm>

Tacon, A.G.J., M.R. Hasan, and R.P. Subasinghe. (2006) Use of fishery resources as feed inputs for aquaculture development: trends and policy implications. *FAO Fisheries Circular* 1018.

Tacon, A.G.J, and I.P. Forster. (2003) *Aquafeeds and the environment: Policy implications.* *Aquaculture*. 226: 181-189

Tacon, A.G.J. (1996) *Feeding Tomorrow's Fish.* World Aquaculture. September 1996.


Trites AW, Miller AJ, Maschner HDG, Alexander MA, Bograd SJ, Calder JA, Capotondi A, Coyle KO, Lorenzo ED, Finney BP, Gregr EJ, Grosch CE, Hare SR, Hunt GL, Jahncke J, Kachel NB, Kim H, Ladd C, Mantua NJ, Marzban C, Maslowski W, Mendelsohn R, Neilson DJ, Okkonen SR, Overland JE, Reedy-Maschner KL, Royer TC, Schwing FB, Wang JXL, Winship AJ. (2007) "Bottom-up forcing and the decline of Stellar sea lions (*Eumetopias jubatus*) in Alaska: assessing the ocean climate hypothesis." *Fish. Oceanography*. 16(1), 46-67. *Jubilation*



- Trites AW, C. V., Pauly D. (2006). Effects of fisheries on ecosystems: just another top predator? Top predators in Marine Ecosystems. Cambridge, Cambridge University Press.
- Trites AW (1998) Stellar Sea Lions (*Eumatopias jubatus*): Causes for their Decline and Factors Limiting their Restoration. Vancouver (BC): Fisheries Centre, University of British Columbia.
- Trites AW, Larkin AP (1992) The Status of Stellar Sea Lion Populations and the Development of Fisheries in the Gulf of Alaska and Aleutian Islands. Vancouver (BC): Fisheries Centre, University of British Columbia.
- Tudela, S. 2005. Risk on local fish populations and ecosystems posed by the use of imported feed fish by the tuna farming industry in the Mediterranean. WWF Mediterranean Program. <http://www.wwf.or.jp/activity/marine/lib/wwfonenvironmentalriskoftunafarming-e.pdf>
- Tuominen, T.R., and M. Esmark. (2003) Food for thought: The use of Marine Resources in Fish Feed. World Wildlife Fund. <http://assets.panda.org/downloads/foodforthoug.pdf>
- Tyedmers, P.H. (2000). Salmon and sustainability: the biophysical cost of producing salmon through the commercial salmon fishery and the intensive salmon aquaculture industry. Ph.D. dissertation, Resource Management and Environmental Studies, University of British Columbia.
- Tynan CT, DeMaster DP. (1997) "Observations and Predictions of Arctic Climatic Change: Potential Effects on Marine Mammals." *Arctic* 50(4), 308-322.
- Uphoff JH Jr. (2003) Predator-prey analysis of striped bass and Atlantic menhaden in Upper Chesapeake Bay 10: 313-322.
- Vaughan, Douglas S., Smith, Joseph W., Williams, Erik H. and Prager, Michael H. (2001) Analysis of the Atlantic Menhaden Stock, NOAA Fisheries, Center for Coastal Fisheries and Habitat Research
- Virginia Institute of Marine Science, Fisheries Science (2003), Research Projects: Menhaden <http://www.fisheries.vims.edu/femap/fish%20pages/Atlantic%20Menhaden.htm>
- Volpe, J.P. Dollars without sense: The bait for big-money tuna ranching around the world. *BioScience* 55(4): 301-302, 2005.
- Walker RV, Myers KW, Davis ND, Aydin KY, Fridland KD, Carlson HR, Boehlert GW, Urawa S, Ueno Y, Anma G. (2000) "Diurnal Variation in the Thermal Environment Experienced by Salmonids in the North Pacific as Indicated by Data Storage Tags" *Fisheries Oceanography* 11, 171-.
- Walter JF III, Overton AS, Ferry KH, Mather ME (2003) Atlantic coast feeding habits of striped bass: a synthesis supporting a coast-wide understanding of trophic biology. *Fisheries Management and Ecology* 10: 349-360
- Wanless S, Wright PJ, Harris MP, Elston DA. (2004) "Evidence for decrease in size of lesser sand eels *Ammodytes marinus* in a North Sea aggregation over a 30-yr period." *Mar. Ecol. Prog. Ser.* 279, 237-246.
- Washington State Department of Ecology, Puget Sound Shoreline, Sand Lance. <http://www.ecy.wa.gov/programs/sea/pugetsound/species/sandlance.html>
- Washington Department of Fish and Wildlife. Forage Fish; Anchovy. <http://wdfw.wa.gov/fish/forage/anchovy.htm>
- Washington Department of Fish and Wildlife, Forage Fish, Sand Lance. <http://wdfw.wa.gov/fish/forage/lance.htm#lbiology>
- Washington State Forage Management Plan: <http://wdfw.wa.gov/fish/forage/manage/foragman.pdf>
- Watson, R., J. Alder, and D. Pauly. (2006) Fisheries for Forage Fish, 1950 to the present. In: Alder J, Pauly D (Ed.) On the multiple uses of forage fish: from ecosystems to markets. Fisheries Centre Research Reports 14(3) Fisheries Centre, University of British Columbia
- Watwood, S.L., P.J.O. Miller, M. Johnson, P.T. Madsen, and P.L. Tyack. (2006) Deep-diving foraging behaviour of sperm whales (*Physeter macrocephalus*). *Journal of Animal Ecology* 75: 814-825.
- Welch, HE; Crawford, RE; Hop H (1993) Occurrence of Arctic cod (*Boreofadus saida*) schools and their vulnerability to predation in the Canadian High Arctic, *Arctic*, Vol. 46, no. 4
- Welch DW, Ishida Y, Nagasawa K. (1998) "Thermal limits and ocean migrations of sockeye salmon long-term consequences of global warming." *Can. J. Fish. Aquat. Sci.* 55, 937-948.
- Whatcom County Marine Resources Committee (2007) Whatcom County MRC Project—Forage Fish. Accessed 5 February 2008. Available at: <http://whatcom-mrc.wsu.edu/MRC/projects/foragefish.htm>
- Whitehead, P.J.P., G.J. Nelson and T. Wongratana, 1988 FAO species catalogue. vo1.7. Clupeoid fishes of the world (Suborder Clupeoidei). An annotated and illustrated catalogue of the herrings, sardines, pilchards, sprats, anchovies and wolf-herrings. Part 2. Engraulididae. For bibliographic purposes this document should be cited as follows: FAO WWF, TRAFFIC North America, 2004 Fisheries Snapshots on Trade and Conservation Issues: Menhaden. http://www.worldwildlife.org/trade/pubs/Menhaden_Snapshot.pdf
- World Wildlife Fund & the Whale and Dolphin Conservation Society. "Whales in Hot Water? The Impact of a Changing Climate on Whales, Dolphins and Porpoises."
- Wright, P., R.T.Barrett, S.P.R. Greenstreet, B.Olsen, and M.L. Tasker. (1996) Effect of fisheries for small fish on seabirds in the eastern Atlantic. In: Hunt, GL, Jr. and RW Furness (eds.) (1996) Seabird/fish interactions, with particular reference to seabirds in the North Sea. ICES Cooperative Research Report No. 216. International Council for the Exploration of the Sea, Copenhagen.
- Zeidberg, D., Louis, The fishery for California market squid (*Loligo opalescens* (Cephalopoda: Myopsida) from 1981 through 2003)monterey bay Aquarium Research institute, UC Marine Council, Coastal Environmental Quality initiative (University of California), paper 032, 2004. <http://repositories.cdlib.org/ucmarine/ceqi/032/>

www.oceana.org/prey





Oceana campaigns to protect and restore the world's oceans. Our team of marine scientists, economists, lawyers and advocates win specific and concrete policy changes to reduce pollution and to prevent the irreversible collapse of fish populations, marine mammals and other sea life. Global in scope and dedicated to conservation, Oceana has campaigners based in North America, Europe, and South America. More than 300,000 members and e-activists in over 150 countries have already joined Oceana. For more information, please visit www.Oceana.org.

