

Ecosystem-based Adaptation in Marine and Coastal Ecosystems



Lynne Zeitlin Hale¹, Imèn Meliane¹, Sarah Davidson¹, Trevor Sandwith¹, Jonathan Hoekstra¹, Mark Spalding¹, Steven Murawski², Ned Cyr², Kenric Osgood², Marea Hatzios³, Pieter Van Eijk⁴, Nicholas Davidson⁵, William Eichbaum⁶, Carlos Dreus⁶, David Obura⁷, Jerker Tamelander⁸, Dorothée Herr⁸, Caleb McClennen⁹, Paul Marshall¹⁰

Climate change is already impacting millions of people, particularly vulnerable communities whose survival, livelihoods and cultural identities are dependent on the integrity of marine and coastal ecosystems. These impacts will continue and increase over the short to medium term, even as the community of nations works to reduce its greenhouse gas emissions. Ecosystem-based adaptation provides an opportunity to reduce the vulnerability of these communities through an improved management of marine and coastal ecosystems so that they continue providing important ecosystem services on which so many depend. There is an urgent need to develop, implement, and fund ecosystem-based adaptation strategies involving coastal communities as a priority response to climate change.

1 The Nature Conservancy

2 National Oceanographic and Atmospheric Administration, USA (NOAA)

3 The World Bank

4 Wetlands International

5 The Ramsar Convention Secretariat

6 World Wide Fund For Nature (WWF)

7 Coastal Oceans Research and Development in the Indian Ocean (CORDIO)

8 International Union for Conservation of Nature (IUCN)

9 Wildlife Conservation Society (WCS)

10 Great Barrier Reef Marine Park Authority, Australia

An initial version of this paper was prepared as a policy brief for The Global Forum on Oceans, Coasts, and Islands for the World Ocean Conference's Global Policy Day (May 13,2009); it was subsequently presented at an Adaptation Workshop held at The International Marine Conservation Congress (May 28,2009). Inputs from participants at both events strengthened the document and are gratefully acknowledged.

Human Societies Depend on Marine and Coastal Ecosystems

The ocean is a unique, extraordinary and vital element of our planet, covering more than 70% of its surface. It sustains life by generating oxygen, absorbing carbon dioxide from the atmosphere, and regulating climate and temperature. Billions of people around the world, especially vulnerable communities in tropical areas, depend on ocean and coastal ecosystems for their survival and well-being. Most of these populations live near (or on) coastlines, and wetlands and reefs provide the first line of coastal defence. More than a billion people worldwide rely on fish as their main source of protein. Fisheries and associated industries employ 38 million people directly, and another 162 million indirectly (FAO, 2008). Nature-based tourism on coral reefs is estimated to contribute \$30 billion to the global economy each year. In addition, marine and coastal ecosystems provide a wide range of other important services to human society, including medicines, natural shoreline protection against storms and floods, water quality maintenance, and other cultural and spiritual benefits (UNEP, 2006).

Coastal ecosystems produce disproportionately more services related to human well-being compared with other systems, even those covering larger total areas, and at the same time they are experiencing some of the most rapid and intense environmental degradation and over-exploitation (Millennium Ecosystem Assessment, 2005).

Climate change is already impacting vulnerable communities that live along coasts

Fifty per cent (50%) of the human population lives along the coast. Population densities in coastal regions are about three times higher than the global average, with **23% of the world's population living both within 100 kilometres of the coast AND less than 100 meters above sea level**. Sixty percent of the world's cities with a population of over 5 million are located within 100 kilometres of the coast. Many of the world's poorest communities also live along the coast and rely on mangrove and reef-based fisheries for food security and on tourism for foreign exchange, particularly in small islands and tropical developing countries. A recent study in Indonesia estimates that 60% of the population is dependent on marine and coastal fishing resources for their protein and livelihoods. In the Wakatobi province, 100% of food requirements are met by the sea, and this is complemented by building materials and cash income derived from marine and coastal natural resources (Emerton, 2009).

Climate change impacts on oceans and coasts are numerous and complex, and expected across polar, temperate, and tropical environments, from the surface to the ocean depths, profoundly altering ecosystem functions (IPCC, 2007; Griffis *et al.*, 2008).

- Rising seas will erode and inundate coastlines and valuable wetlands and can increase salinity in coastal water sources and lands used to produce food.
- Increased water temperatures make corals more vulnerable to bleaching and change the geographical ranges of many marine species. Already 20% of the world's coral reefs are estimated to be damaged beyond repair and unless emissions are drastically reduced, 80% of the world's reefs could be lost, and their ability to feed people and sustain the livelihoods of millions of people severely impaired.
- Increasing acidification of the oceans as a result of CO₂ absorption reduces the ability of key marine organisms like corals, plankton, and shellfish to build shells and skeletons, with consequences for the productivity of marine ecosystems and dependent fisheries.
- Changing weather, wind patterns and sea temperatures impact various oceanographic processes, including nutrient upwellings and surface currents, changing population abundance and distribution for many marine species, affecting ocean productivity.
- Predicted decline in oxygen concentrations across various depths, reduced ventilation of the mid-water from ocean warming, and local eutrophication will lead to an expansion of oceanic dead zones and fish mortality.

Climate change is also exacerbating other threats to the oceans in a cumulative way. The increased occurrence and distribution of invasive alien species, growing pressure on fisheries resources, altered patterns of coastal development and shore-based pollution are all exacerbated by climate change leading to increased vulnerability and an unpredictable scale of impact.

The Global Response to Climate Change – Mitigation and Adaptation

There is an immediate need for a significant reduction in greenhouse gas emissions to reduce the rate of climate change and to avoid catastrophic impacts on biodiversity in the long term. In the absence of such strong mitigation action, it is possible that the most vulnerable ecosystems, such as coral reefs, will cease to function in their current forms within a few decades (Hoegh-Guldberg *et al.*, 2007).

Even if mitigation measures are effectively implemented, the earth's climate will continue changing over the short to medium term, due to lag effects of temperature and ocean acidification in response to the build-up of CO² already in the atmosphere. This will lead to unavoidable impacts requiring adaptive management responses in the face of change. Management actions can exacerbate or ameliorate the situation. Adaptive measures to reduce impacts and to increase resilience in the face of these changes are a necessary complement to mitigation actions, and will include measures to ensure ecosystem integrity, reduce manageable impacts like pollution, restore habitats, alter use patterns, and most important avoid inadvertent measures that address one problem, like coastal inundation, but cause others, e.g. destruction of coastal ecosystems. Adaptation measures will also increasingly employ the regenerative capacity of natural ecosystems as part of engineered solutions.

There is a need for comprehensive adaptation strategies to consider not only “hard infrastructure” but also ecosystem-based solutions

Coastlines are now more dynamic than ever because of changing storm patterns and sea level rise, placing human and natural communities at greater risk. The costs of these hazards to human and natural communities are increasing as coastal development continues and natural buffers, such as coastal wetlands and dunes, are lost. One of the areas where there are real opportunities for identifying win-win solutions for human and natural communities is in building approaches that combine hazard mitigation and biodiversity conservation in coastal zones to preserve infrastructure, protect human communities and preserve their livelihoods (Kareiva and Marvier, 2007).

Most existing and proposed adaptive responses to climate change in coastal areas have focused on using “hard” engineering solutions to try to address the problem. Over the past century, hard coastal defence structures have become ubiquitous features of coastal landscapes as a response to these threats. The proliferation of defence works affects over half of the shoreline in some regions and results in dramatic changes to the coastal environment. The extent and projected trend of this phenomenon are alarming. For example, 22000 km² of the coastal zone in Europe are covered in concrete or asphalt, and urban artificial surfaces increased by nearly 1900 km² between 1990-2000 alone (Airoldi and Beck, 2007). Similar examples occur in other parts of the world - e.g. California, Australia and Japan - where hundreds of kilometres of coasts are hardened. Worse, the addition of one artificial structure changes erosion and sediment transport and creates the need for another structure just down-shore in a costly negative feedback loop (Airoldi *et al.*, 2005).

Such expensive infrastructural responses, though in some cases necessary, will not be sufficient to address the full scope of climate change impacts, and can exacerbate the destruction of fragile ecosystems, further reducing their ability to adapt. For example, shoreline hardening adversely affects wetlands through direct destruction and by preventing sediment transport essential to that ecosystem. This results in increased erosion and further loss of habitat on directly adjacent or downstream shorelines (US EPA, 2009). In addition, such options often come with high maintenance costs.

Ecosystem-based adaptation options often apply directly or indirectly to multiple management goals. For example, allowing wetlands to migrate inland will not only maintain their shoreline protection services, but could also directly address maintaining water quality and preserving habitat for maintaining local fisheries or tourism. Managers need to take into consideration priorities, costs and trade-offs and consider implementing different options in different areas according to which resources are most in need of protection. In some cases, integrating “soft” and “hard” engineering approaches to adaptation could allow for the development of structural measures targeted at protecting the natural ecosystems themselves, in cases where climate impacts extend beyond their natural resilience. In the Mississippi Delta, for example, plans are being developed for the construction of small dikes that protect salt marshes and coastal peatlands against erosion and allow them to naturally regenerate. Subsequently, the regenerated coastal ecosystems contribute to the resilience of the Delta as a whole and are able to provide their full range of services.

Ecosystem-based Adaptation

Ecosystem-based adaptation aims to:

- Preserve and restore natural ecosystems that can provide cost-effective protection against some of the threats that result from climate change. For example, coastal ecosystems like wetlands, mangroves, coral reefs, oyster reefs, and barrier beaches all provide natural shoreline protection from storms and flooding in addition to their many other services (CBD, 2009).
- Conserve biodiversity and make ecosystems more resistant and resilient in the face of climate change so that they can continue to provide the full suite of natural services. This is particularly important for sustaining natural resources (e.g., fish stocks, fuel, clean water, marine biodiversity for tourism attractions) on which vulnerable communities depend for their subsistence and livelihoods.

Ecosystem-based adaptation requires collective action among governments, communities, conservation and development organisations, and other stakeholders to plan and empower action that will enhance environmental and community resilience to climate change impacts. In addition, it can be a major opportunity for community-based adaptation. Vulnerable coastal communities can be engaged, employ local knowledge and participate directly in developing and applying ecosystem-based solutions.

Ecosystem-based Adaptation Benefits in Marine and Coastal Areas

Adaptation strategies that aim to enhance the resilience of ecosystems to enable the continued provision of goods and services can be particularly important for vulnerable communities, who are often directly dependent upon natural resources. A growing body of evidence suggests that ecosystem-based adaptation can be a cost-effective strategy across the major adaptation sectors (Campbell *et al.*, 2009). In addition, ecosystem-based adaptation strategies often address multiple coastal management goals and provide multiple benefits (see Table 1; US EPA, 2009).

Below are some examples of benefits of ecosystem-based adaptation strategies.

Cost-effective shoreline protection. Hard infrastructure like seawalls and levees is expensive, requires ongoing maintenance, and can fail catastrophically under severe storm conditions. Alternatively, an ecosystem-based approach of protecting and restoring “green infrastructure” like healthy coastal wetlands, including mangrove forests and coral reefs, could be a more cost-effective means of protecting large coastal areas, requiring less maintenance (Moberg and Rönnbäck, 2003). Across the globe, there are numerous examples of the important role that coastal ecosystems such as mangroves, wetlands, shellfish reefs and coral reefs play in coastal protection as they dissipate wave energy. Mangrove restoration in Vietnam has been shown to attenuate wave height and thus reduce wave damage and

erosion (Mazda *et al.*, 1997). Sri Lanka's Muthurajawela marsh, a coastal peat bog covering some 3100 hectares, is an important part of local flood control. In Malaysia, the value of intact mangrove swamps for storm protection and flood control has been estimated at US \$300,000 per km, which is the cost of replacing them with rock walls (Ramsar Convention on Wetlands, 2005). Healthy mangroves also provide numerous additional benefits, such as timber and fisheries production, biofiltration, and recreational activities like recreational fishing and bird watching (Spalding *et al.*, in press), services not provided by non-ecosystem-based coastal protection alternatives.

Shellfish reefs serve as natural coastal buffers, absorbing wave energy directed at shorelines and reducing erosion from boat wakes, sea level rise, and storms (Meyer 1997, Piazza *et al.* 2005). In addition, shellfish reefs play an important role as habitat for other species; the fish produced on oyster reefs have significant value to coastal economies (Beck *et al.*, 2009; Grabowski and Peterson, 2007). They also improve water quality through filtration and provide fish habitat, which can enhance tourism and recreation (Freeman, 1995; Lipton 2004).

The analysis of recent disasters, such as the December 2004 Indian Ocean tsunami and the hurricanes that struck North and Central America in September and October 2005, demonstrate the importance of habitat protection and natural resource management in decreasing vulnerability to extreme events (Sudmeier-Rieux *et al.*, 2006). Prominent insurers and investors are likewise incorporating and advocating risk reduction using the protective value of ecosystems and other natural infrastructure, such as coastal wetlands, barrier islands, trees, mangroves and other vegetation, as part of development appraisals. This reflects the industry's understanding that natural infrastructure is essential to society's efforts to address climate change, and that these systems must be included as part of any adaptation strategy (Heinz Center and Ceres, 2009).

Mangroves, saltmarshes and seagrasses are critically important sediment traps while their high productivity enables them to add considerable volume to trapped sediments around their roots. The result is that their soils can "grow" upwards, enabling them to keep pace with rising sea levels or at least reduce the relative rate of sea level rise compared to unvegetated sites. In some mangrove locations in the Caribbean, mangrove sediments are rising at over 4mm/year (Cahoon and Lynch, 1997; McKee *et al.*, 2007), significantly above the recent global mean sea level rise of 3.1mm/year.

Sustaining local livelihoods and contributing to local economies. The World Bank's Climate Change Framework Strategy (2008) warns that the disproportionate impacts of climate change on the poorest and most vulnerable communities could set back much of the development progress of the past decades and plunge communities back into poverty. Ecosystem-based adaptation helps maintain ecosystem productivity and supports sustainable income-generating activities in the face of climate change. For example, in Kimbe Bay, Papua New Guinea, coral reef resilience principles were applied to design a network of marine protected areas to help the Bay's ecosystems withstand the impacts of a warming ocean and continue to provide food and other resources to local communities (Green *et al.*, 2009). In Samoa, mangroves are being planted as part of a larger restoration project to enhance food security and protect local communities from storm surges, which are expected to increase as a result of climate change (UNDP, 2008). In Myanmar, communities are replanting mangroves in the Ayerwaddy Delta following the destruction from Cyclone Nargis, which devastated life and property in the absence of mangrove forests that had been cleared over time for paddy cultivation (Tripartite Core Group, 2008).

The contribution of marine and coastal ecosystem services to local, regional, and national economies is substantial. For example, a recent study (Emerton, 2009) of the value of Indonesia's coastal ecosystems identified a potential value of sustainable fisheries from coral reef areas alone of more than US\$1.2 billion—almost half of the value of national fisheries production. The same study found that marine and coastal ecosystems are responsible for about 49% of the Keladupa sub-district economy, and coral reef fisheries provide the main source of income for almost 80% of the residents in the Raja Ampat Regency. Marketed mangrove products generate 22% of the local economy of Ranong province in Thailand (IUCN, 2008). Estimates of direct tourism revenue generated from the presence and use of medium to good quality coral reefs in the Philippines range from US \$38,000 to \$63,000 per km² (White and Trinidad, 1998).

Carbon sequestration and reinforcement of mitigation efforts. Coastal wetlands, including marshes and mangroves, sequester substantial amounts of carbon (Pritchard, 2009), so also play a crucial and incremental role in reducing the pace and scale of climate change itself. Mangroves may play an important role in carbon sequestration and storage in local and *ex situ* sediments; around 10% of mangrove productivity is incorporated into local sediments (Spalding *et al.*, in press). Other studies have estimated that mangroves contribute about 15% of the organic carbon accumulating in marine sediments globally (Twilley *et al.* 1992; Jennerjahn and Ittekkot 2002). A conservative estimate is that mangroves sequester an estimated 112 ± 85 Tg C per year, which is still an underestimation due to the lack of information about fine root activities. This amount of carbon sequestration is comparable with that for tropical terrestrial forests (Alongi, 2008; Bouillon *et al.*, 2007). The majority of this captured carbon is likely to remain stable over millennial time-scales, making mangroves an important carbon “sink”.

Overall, it has recently been estimated that marine angiosperms (including saltmarsh, mangrove and seagrass) contribute some 46% of total organic carbon buried into marine sediments, or some 111 teragrams per year (Spalding *et al.*, in press). This contribution by coastal vegetation had been overlooked in earlier mass balance studies and such inputs represent a near-doubling of earlier estimates of the carbon storage function of marine sediments, making them highly significant contributors to global models of carbon flows (Duarte *et al.* 2005).

Providing refugia - a place to hide. Marine protected areas (MPAs) act as refugia, protecting critical areas and functions in the life cycles of important marine species (IUCN-WCPA, 2008). Refugia are important to protect species and larval sources which aid in the recovery of damaged areas. Well-designed MPAs and MPA networks have proven to be an important tool in increasing the resilience and adaptive capacity of coral reefs to bleaching, by protecting them from other disturbances such as increased nutrient loads, pollution, diver and boat damage, sedimentation, and destructive and overfishing (Smith *et al.*, 2009). Existing research and management practices have demonstrated that connectivity among sites within an MPA network helps insure against the risk of losing an important habitat or community type following a disturbance such as a bleaching episode or intense storm. The widespread replication of these experiences for increasing the resilience of MPA networks in the face of climate change impacts provides a solid foundation for rapid expansion of these important management approaches as a key strategy for protecting ocean and coastal ecosystem services and the wide range of benefits they provide us.

Contributing to social resilience. Communities and local decision-makers still have little access to information on likely changes that will impact their lives and livelihoods and to tools to visualise the potential impacts and identify alternative scenarios. As a consequence, communities are unable to integrate climate-change related impacts and risks into decision-making regarding natural resource protection and land use management. Development of such tools is beginning to occur on pilot scales, [e.g. the Cristal assessment tool (<http://www.cristaltool.org/>), the assessment of livelihood vulnerability and adaptation options for communities dependent on coral reefs (Marshall *et al.*, 2009)], but much more work is needed.

Ecosystem-based adaptation provides a major opportunity for community-based adaptation. By maintaining and restoring healthy ecosystems that are more resilient to climate change impacts, ecosystem-based strategies can help ensure continued availability and access to water and other essential natural resources and ecosystem services so that vulnerable communities can better cope with climate variability and change. These communities can be engaged, employ local knowledge and participate directly in developing and applying ecosystem-based solutions.

Table 1. Adaptation options for maintaining and restoring coastal wetlands and shorelines. (Adapted from US EPA, 2009)

Adaptation Option	Climate Stressor Addressed	Additional Management Goals Addressed	Benefits	Constraints
Allow coastal wetlands to migrate inland (e.g., through setbacks, density restrictions, land purchases)	Sea level rise	Preserve habitat for vulnerable species; Preserve coastal land/development	Maintains species habitats; maintains protection for inland ecosystems	In highly developed areas, there is often no land available for wetlands to migrate, or it can be costly to landowners
Incorporate wetland protection into infrastructure planning (e.g., transportation planning, sewer utilities)	Sea level rise; Changes in precipitation	Maintain water quality; Preserve habitat for vulnerable species	Protects valuable and important infrastructure	
Preserve and restore the structural complexity and biodiversity of vegetation in tidal marshes, seagrass meadows, and mangroves	Increases in water temperatures; Changes in precipitation	Maintain water quality; Maintain shorelines; Invasive species management	Vegetation protects against erosion, protects mainland shorelines from tidal energy, storm surge, and wave forces, filters pollutants, and absorbs atmospheric CO ₂	
Identify and protect ecologically significant ("critical") areas such as nursery grounds, spawning grounds, and areas of high species diversity	Altered timing of seasonal changes; Increases in air and water temperatures	Invasive species management; Preserve habitat for vulnerable species	Protecting critical areas will promote biodiversity and ecosystem services (e.g., producing and adding nutrients to coastal systems, serving as refuges and nurseries for species)	May require federal or state protection
Integrated Coastal Zone Management (ICZM) – using an integrated approach to achieve sustainability	Changes in precipitation; Sea level rise; Increases in air and water temperatures; Changes in storm intensity	Preserve habitat for vulnerable species; Maintain/restore wetlands; Maintain water availability; Maintain water quality; Maintain sediment transport; Maintain shorelines	Considers all stakeholders in planning, balancing objectives; addresses all aspects of climate change	Stakeholders must be willing to compromise; requires much more effort in planning
Incorporate consideration of climate change impacts into planning for new infrastructure (e.g., homes, businesses)	Sea level rise; Changes in precipitation; Changes in storm intensity	Preserve habitat for vulnerable species; Maintain/restore wetlands	Engineering could be modified to account for changes in precipitation or seasonal timing of flows; siting decisions could take into account sea level rise	Land owners will likely resist relocating away from prime coastal locations
Create marsh by planting the appropriate species – typically grasses, sedges, or rushes – in the existing substrate	Sea level rise	Maintain water quality; Maintain/restore wetlands; Preserve habitat for vulnerable species; Invasive species management	Provides protective barrier; maintains and often increases habitat	Conditions must be right for marsh to survive (e.g., sunlight for grasses, calm water); can be affected by seasonal changes
Use natural breakwaters of oysters (or install other natural breakwaters) to dissipate wave action and protect shorelines	Increases in water temperatures; Sea level rise; Changes in precipitation; Changes in storm intensity	Preserve coastal land/development; Maintain water quality; Invasive species management	Naturally protect shorelines and marshes and inhibit erosion inshore of the reef; will induce sediment deposition	May not be sustainable in the long-term, because breakwaters are not likely to provide reliable protection against erosion in major storms
Replace shoreline armoring with living shorelines – through beach nourishment, planting vegetation, etc.	Sea level rise; Changes in storm intensity	Maintain/restore wetlands; Preserve habitat for vulnerable species; Preserve coastal land/development	Reduces negative effects of armoring (down-drift erosion); maintains beach habitat	Can be costly; requires more planning and materials than armoring
Remove shoreline hardening structures such as bulkheads, dikes, and other engineered structures to allow for shoreline migration	Sea level rise	Maintain sediment transport	Allows for shoreline migration	Costly for, and destructive to, shoreline property
Plant SAV (such as sea grasses) to stabilize sediment and reduce erosion	Changes in precipitation; Sea level rise	Maintain/restore wetlands; Preserve habitat for vulnerable species; Preserve coastal land/development	Stabilizes sediment; does not require costly construction procedures	Seasonality – grasses diminish in winter months, when wave activity is often more severe because of storms; light availability is essential

GUIDING PRINCIPLES AND RECOMMENDATIONS FOR ECOSYSTEM-BASED ADAPTATION

Guiding principles for developing effective ecosystem-based adaptation strategies include:

- ***Nature's infrastructure should be used first.*** Natural ecosystems provide valuable protection and other services, and we should take advantage of them. Maintaining and restoring "nature's infrastructure" should be a priority for reducing vulnerability to climate change impacts. As the effects of climate change become more severe, there will be, however, situations where engineering and hard structures may be necessary. Such structures need be built in sync with nature and its changing patterns.
- ***Healthy ecosystems will be more resilient to climate changes.*** Ecosystem-based adaptation strategies should include a focus on minimising other anthropogenic stresses that have degraded the condition of critical ecosystems. It is also important to take into account the full range of impacts, as one environmental change may have cascading effects.
- ***Existing management practices and governance infrastructure should be improved.*** The most effective ecosystem-based strategies currently available apply established best practices in land, water, and natural resource management to confront the new challenges posed by climate change. **Effective management programs** that address multiple stressors and that take into consideration priorities, trade-offs and synergies are central to adaptation planning. Well-designed and effectively managed **marine protected area networks** can make an enormous contribution to maintaining natural connections across seascapes so that ecosystems can continue to function and to provide services to dependent communities (Smith *et al.*, 2009).
- ***Stakeholders should be involved in strategy development.*** Ecosystem-based adaptation presents a tangible opportunity to solve climate change problems by aligning conservation, development, and poverty alleviation interests. Such synergies benefit from government collaboration with indigenous and local communities, conservationists, relevant private sector stakeholders, development specialists, and humanitarian aid specialists.
- ***Decision support tools help visualise future scenarios and compare alternative adaptations.*** One of the major impediments to decision-making is a visceral understanding of potential impacts from climate change to communities and their resources. Tools that can help visualise these futures can be as simple as pictures of coastlines with different flooding scenarios from sea level rise and storms to interactive map servers. They also can provide the basis for examining costs and benefits of alternative approaches to adaptation with either hard or soft solutions with a goal of reducing losses for human and natural communities.
- ***Government and the private sector can provide incentives for "climate smart" development and discourage development in vulnerable and sensitive habitats.*** The financial and insurance sectors can and need to play a positive role in ecosystem-based adaptation by fully recognising and accounting for risks associated with development in vulnerable areas and providing incentives for maintaining "nature's infrastructure."
- ***Environmental, ecological, social and economic changes should be measured and mapped.*** As climate change impacts increase and our scientific understanding and observation evolve over time, it is important to monitor and report these changes and build on them to improve predictions, and to adapt responses.

- **Adaptive management is imperative.** While the general trends in climate change are well-documented, the timing and magnitude of local changes remain difficult to predict accurately. Ecosystem-based adaptation strategies should include monitoring so that management actions can be quickly adjusted in response to changing conditions. Management objectives may need to be revised and geographic priorities may need to be reconsidered to protect natural climate change “refugia”, or to triage places suffering severe climate change impacts.
- **“Mainstreaming” ecosystem-based adaptation into coastal management and development at all levels.** Ecosystem-based strategies will be more effective if they are mainstreamed into other development initiatives such as poverty reduction strategies, country development strategies and sector plans and “owned” by those authorities responsible for preparing and implementing them.
- **A regional approach is needed.** Ecosystems stretch beyond political and geographical boundaries, and this is particularly true for the marine environment. Therefore, efforts need to be made to design adaptation measures that are not limited by these boundaries. Adaptation measures for a resource shared by multiple states can succeed only through integration of a regional or transboundary dimension.
- **Prepare for the unexpected.** In preparing for climate change, we need to keep in mind the possibility of non-linear, abrupt changes or step functions which can alter the state of an ecosystem or biome quickly once a threshold has been reached. These uncertain but high consequence events (such as de-glaciation or alteration of oceanic currents) need to be acknowledged and social resilience to cope with such changes developed.

Currently, a growing number of local, national and regional initiatives and projects are applying ecosystem-based adaptation principles to a variety of marine and coastal areas and in various parts of the globe (e.g., see Table 1). There is a need to synthesise the new results as they become available, develop additional management tools and transfer technology and build capacity for their use.

References

- Airoldi, L., M. Abbiati, M. W. Beck, S.J. Hawkins, P.R. Jonsson, D. Martin, P. S. Moschella, A. Sundelöf, R. C. Thompson, P. Åberg. 2005. An ecological perspective on the deployment and design of low- crested and other hard coastal defence structures. *Coastal Engineering* 52: 1073–1087.
- Airoldi, L., M. W. Beck. 2007. Loss, status, and trends for coastal marine habitats of Europe. *Annual Review of Marine Biology and Oceanography* 45:347-407.
- Alongi, D.M. 2008. Mangrove forests: Resilience, protection from tsunamis, and responses to global climate change. *Estuarine, Coastal and Shelf Science*, 76, 1-13.
- Beck, M. W., R.D. Brumbaugh, L. Airoldi, A. Caranza, L. D. Coen, C. Crawford, O. Defeo, G.J. Edgar, B.Hancock, M. Kay, H. Lenihan, M.W. Luckenbach, C.L. Toropova, G. Zhang. 2009. Shellfish Reefs at Risk: A Global Analysis of Problems and Solutions. The Nature Conservancy, Arlington VA. 52 pp.
- Bouillon, S., A. V. Borges, E. Castañeda-Moya, K. Diele, T. Dittmar, N. C. Duke, E. Kristensen S. Y. Lee, C. Marchand, J. J. Middelburg, V. H. Rivera-Monroy, T. J. Smith III, R. R. Twilley. 2008. Mangrove production and carbon sinks: A revision of global budget estimates, *Global Biogeochem. Cycles*, 22, GB2013.
- Cahoon DR, Lynch JC (1997) Vertical accretion and shallow subsidence in a mangrove forest of southwestern Florida, U.S.A. *Mangroves and Salt Marshes* 1: 173
- Campbell A., V. Kapos, A. Chenery, S.I. Kahn, M. Rashid, J.P.W. Scharlemann, B. Dickson . 2009. The linkages between biodiversity and climate change mitigation UNEP World Conservation Monitoring Centre.
- CBD. 2009. *Identifying and Enhancing the Linkages between Biodiversity and Climate-Change Adaptation*. UNEP/CBD/AHTEG/BD-CC-2/2/3
- Duarte, C.M., J. J. Middelburg, N. Caraco.2005. Major role of marine vegetation on the oceanic carbon cycle. *Biogeosciences* 2: 1-8
- Emerton, L. 2009. Investing in natural infrastructure: the economic value of Indonesia’s marine protected areas and coastal ecosystems. Unpublished report. The Nature Conservancy.
- FAO. 2008. *The State of World Fisheries and Aquaculture 2008*. Food and Agriculture Organization, Rome.

- Freeman, M. 1995. The benefits of water quality improvements for marine recreation: a review of the empirical evidence. *Marine Resource Economics* 10:385-406.
- Grabowski, J. H., C. H. Peterson. 2007. Restoring oyster reefs to recover ecosystem services. Pages 281-298 in K. Cuddington, J. Byers, W. Wilson, and A. Hastings, editors. *Ecosystem engineers: plants to protists*. Academic Press, Boston.
- Green, A., S.E. Smith, G. Lipsett-Moore, C. Groves, N. Peterson, S. Sheppard, P. Lokani, R. Hamilton, J. Almany, J. Aitsi and L. Bualia. 2009. Designing a Resilient Network of Marine Protected Areas for Kimbe Bay, Papua New Guinea. *Oryx* 96:2 (in press)
- Griffis, R.B., R.L. Feldman, N.K. Beller-Simms, K.E. Osgood, and N. Cyr (eds). 2008. *Incorporating climate change into NOAA's stewardship responsibilities for living marine resources and coastal ecosystems: A strategy for progress*. U.S. Dep. Commerce, NOAA Tech. Memo. NMFS-F/SPO-95, 89 p.
- Heinz Center and Ceres. 2009. *Resilient Coasts: A Blueprint for Action*.
- Hoegh-Guldberg, O., P. J. Mumby, A. J. Hooten, R. S. Steneck, P. Greenfield, E. Gomez, C. D. Harvell, P. F. Sale, A. J. Edwards, K. Caldeira, N. Knowlton, C. M. Eakin, R. Iglesias-Prieto, N. Muthiga, R. H. Bradbury, A. Dubi, M. E. Hatzitolos. 2007. Coral reefs under rapid climate change and ocean acidification. *Science*, Vol. 318. no. 5857, pp. 1737 - 1742
- IPCC. 2007. *Climate Change 2007: Synthesis Report*. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change. Core Writing Team, Pachauri, R.K and Reisinger, A. (eds.). IPCC, Geneva, Switzerland.
- IUCN. 2008. Ecological and socio-economic values of Mangrove ecosystems in tsunami affected areas: Rapid ecological-economic-livelihood assessment of Ban Naca and Ban Bangman in Ranong Province, Thailand. IUCN publication. 30pp.
- IUCN World Commission on Protected Areas (IUCN-WCPA). 2008. Establishing Marine Protected Area Networks—Making It Happen. Washington, D.C.: IUCN-WCPA, National Oceanic and Atmospheric Administration and The Nature Conservancy. 118 p.
- Jennerjahn, T., V. Ittekkot. 2002. Relevance of mangroves for the production and deposition of organic matter along tropical continental margins. *Naturwissenschaften* 89: 23
- Kareiva, P., M. Marvier. 2007. Conservation for the People. *Scientific American* 2007:50-57.
- Lipton, D. 2004. The value of improved water quality to Chesapeake Bay boaters. *Marine Resource Economics* 19:265-270.
- Marshall, N.A., P.A. Marshall, J. Tamelander. 2009, in prep. Sustaining Tropical Coastal Communities & Industries: A Framework for Social Adaptation to Climate Change. IUCN Climate change and Coral Reefs working group. www.iucn.org/cccr
- Mazda, Y., M. Magi, M. Kogo and P.N. Hong. 1997. Mangroves as a coastal protection from waves in the Tong King delta, Vietnam. *Mangroves and Salt Marshes*. 1(2): 127-135.
- McKee KL, Cahoon D, Feller IC (2007) Caribbean mangroves adjust to rising sea-level through biotic controls on soil elevation change. *Global Ecology and Biogeography* DOI: 10.1111/j.1466-8238.2007.00317.x: 12
- Meyer, B. L. 1997. Stabilization and erosion control value of oyster cultch for intertidal marsh. *Restoration Ecology* 5:93-99.
- Millennium Ecosystem Assessment. 2005. www.millenniumassessment.org.
- Moberg, F., P. Ronnback. 2003. Ecosystem services of the tropical seascape: Interactions, substitutions and restoration, *Ocean and Coastal Management* 46. pp. 27–46.
- Piazza, B. P., P. D. Banks, and M. K. La Peyre. 2005. The potential for created oyster shell reefs as a sustainable shoreline protection strategy in Louisiana. *Restoration Ecology* 13:499-506.
- Pritchard, D. 2009. *Reducing Emissions from Deforestation and Forest Degradation in developing countries (REDD) – the link with wetlands*. Foundation for International Environmental Law and Development, London.
- Ramsar Convention on Wetlands. 2005. www.ramsar.org/values_shoreline_e.htm.
- Smith, S E., I. Meliane, A. White, C. Snyder, B. Cicin-Sain, and R. Danovaro 2009. *Impacts of Climate Change on Marine Biodiversity and the Role of Networks of Marine Protected Areas*, Policy Brief 7a prepared by Global Forum on Oceans, Coasts and Islands for the World Ocean Conference.
- Spalding, M.D., M. Kainuma, L. Collins. In press. *World Mangrove Atlas*. Earthscan, with International Society for Mangrove Ecosystems, Food and Agriculture Organization of the United Nations, The Nature Conservancy, UNEP World Conservation Monitoring Centre, United Nations Scientific and Cultural Organisation, United Nations University, London
- Sudmeier-Rieux, K., H. Masundire, A. Rizvi, S. Rietbergen (eds). 2006. *Ecosystems, Livelihoods and Disasters: An integrated approach to disaster risk management*. IUCN, Gland, Switzerland and Cambridge, UK. x + 58 pp.
- Tripartite Core Group. 2008. *Post Nargis Joint Assessment*. Government of the Union of Myanmar, United Nations, ASEAN.
- Twilley, R.R., R.H. Chen, T. Hargis. 1992. Carbon sinks in mangroves and their implications to carbon budget of tropical coastal ecosystems. *Water, Air, and Soil Pollution* 64: 265-288
- UNDP. 2008. *Community Based Adaptation: Samoa*. United Nations Development Programme

UNEP. 2006. *Marine and coastal ecosystems and human wellbeing: A synthesis report based on the findings of the Millennium Ecosystem Assessment*. United Nations Environment Programme

US. EPA. 2009. Synthesis of adaptation options for coastal areas. Washington DC, Us Environmental Protection Agency, climate ready estuaries program. EPA 430-F-08-024, January 2009.

White, A.T., A. Cruz-Trinidad. 1998. The Values of Philippine Coastal Resources: Why Protection and Management are Critical. Coastal Resource Management Project, Cebu City, Philippines, 96p.

World Bank. 2008. *Development and Climate Change: A Strategic Framework for the World Bank Group*. World Bank, Washington, DC. Available at: www.worldbank.org/climatechange



Besides providing important coastal protection, fuel and food sources, mangroves and other coastal ecosystems host a range of biodiversity that maintains important tourism activities and local economies.

Photo Credits:

Front Page, from left to right:

Aerial view of Palau. Imen Meliane

Coral reefs of Fiji. Daniel & Robbie Wisdom

Fishermen in Ecuador. Imen Meliane

This Page:

Sian Ka'an Biosphere Reserve, Mexico. Lynn Mc Bride