

# **Green Turtle Nesting Sites and Sea Turtle Legislation throughout Oceania**

Kimberly A. Maison,  
Irene Kinan-Kelly, and  
Karen P. Frutchey



U.S. Department of Commerce  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service

NOAA Technical Memorandum NMFS-F/SPO-110  
September 2010



# Green Turtle Nesting Sites and Sea Turtle Legislation throughout Oceania

Kimberly A. Maison and Karen P. Frutchey  
Joint Institute for Marine and Atmospheric Research  
University of Hawaii  
1000 Pope Road  
Honolulu, Hawaii 96822, USA

Irene Kinan-Kelly  
National Marine Fisheries Service  
Pacific Islands Regional Office  
Protected Resources Division  
1601 Kapiolani, Blvd. #1110  
Honolulu, Hawaii 96814, USA

**NOAA Technical Memorandum NMFS-F/SPO-110**  
**September 2010**



U.S. Department of Commerce  
Gary Locke, Secretary of Commerce

National Oceanic and Atmospheric Administration  
Jane Lubchenco, Ph.D., Administrator of NOAA

National Marine Fisheries Service  
Eric C. Schwaab, Assistant Administrator for Fisheries

**Suggested citation:**

Maison, K.A., Kinan-Kelly, I. and K.P. Frutchey. 2010. Green Turtle Nesting Sites and Sea Turtle Legislation throughout Oceania. U.S. Dep. Commerce, NOAA Technical Memorandum. NMFS-F/SPO-110, 52 pp.

**A copy of this report may be obtained from:**

National Marine Fisheries Service  
Pacific Islands Regional Office  
Protected Resources Division  
1601 Kapiolani, Blvd. #1110  
Honolulu, Hawaii 96814  
Irene.Kelly@noaa.gov

**Or online at:**

<http://spo.nmfs.noaa.gov/tm/110.pdf>

## **Summary**

This is a literature review compiled by the National Marine Fisheries Service (NMFS) Pacific Islands Regional Office (PIRO) Protected Resources Division (PRD) to facilitate better understanding of green turtle nesting distribution in the Central and Western Pacific Ocean, including a summary of legal protections for sea turtles throughout the region. This synthesis of 189 Pacific green turtle nesting locations, nesting assemblage characteristics where available, and current national legislation provides the background information necessary to identify where gaps exist in monitoring and management. This will help NMFS (or other relevant stakeholders) prioritize and direct future research, management activities, and international collaborations to advance green turtle conservation and recovery efforts in the Pacific.



## Table of Contents

Summary.....	iii
Table of Contents.....	v
List of Tables and Figures.....	vi
Introduction.....	1
Methods.....	2
Synthesis of Existing Data on Green Turtle Nesting and Legislation in Oceania.....	4
U.S. States, Territories, and Possessions.....	4
American Samoa.....	4
Commonwealth of the Northern Mariana Islands.....	5
Guam.....	6
Hawaii.....	7
Pacific Remote Island Areas.....	8
International – States with Compacts of Free Association.....	9
Federated States of Micronesia.....	9
Republic of the Marshall Islands.....	12
Republic of Palau.....	13
International.....	14
Australia.....	14
Cook Islands.....	17
Fiji.....	18
French Polynesia.....	18
Japan.....	19
Kiribati.....	20
Nauru.....	21
New Caledonia.....	21
Niue.....	22
Papua New Guinea.....	22
Pitcairn Islands.....	24
Samoa.....	24
Solomon Islands.....	25
Tokelau.....	26
Tonga.....	26
Tuvalu.....	27
Vanuatu.....	27
Wallis and Futuna.....	28
Discussion and Recommendations.....	29
Acknowledgments.....	33
Tables and Figures.....	35
Appendix I: Pacific Countries and Existing International Arrangements.....	41
References.....	43

## List of Tables and Figures

Table 1.	Summary of information for nesting green turtles in Oceania including current generalized legislation .....	35
Figure 1.	Nesting sites for green turtles throughout Oceania and the estimated minimum annual nesting females at each site along with categories of legislation associated with each country.....	37
Figure 1a.	Green turtle nesting and legislation in Oceania: Micronesia/Melanesia...	38
Figure 1b.	Green turtle nesting and legislation in Oceania: Polynesia.....	38
Figure 2.	Locations of active monitoring projects at green turtle nesting beaches throughout Oceania that have collected data within the past five years (since 2005).....	39



# Green Turtle Nesting Sites and Sea Turtle Legislation throughout Oceania

## Introduction

Green turtles (*Chelonia mydas*) in the Pacific Ocean are widely distributed, nesting at hundreds of sites among thousands of islands and atolls scattered throughout the Western and Central Pacific Ocean (WCPO or Oceania). Green turtles are listed as *threatened* under the U.S. Endangered Species Act (ESA) and *endangered* under the internationally recognized IUCN Red List of Threatened Species. They are protected under varying degrees of national legislation and international arrangements throughout the region. For the purposes of this document, Oceania refers to Polynesia, Micronesia and Melanesia extending from Hawaii west to Japan, south to Australia, and southeast to the Pitcairn Islands—a total of 22 sovereign nations, including the U.S. which is represented by the State of Hawaii, three U.S. Territories and seven Pacific Remote Island Areas (PRIAs) under U.S. possession. The quality and amount of green turtle nesting information available is variable among countries in this region as some sites have active monitoring projects and recent data whereas others are not actively monitored. Overall, little information exists regarding stock structure, abundance, or trends for green turtles in this vast region. However, a range of nesting records and survey information is available in published documents, gray literature, and in-country government reports gathered over the past 30+ years representing varying degrees of effort and consistency in monitoring.

There have been prior efforts to compile sea turtle nesting information from this region (Bjorndal 1982 and revised in 1995; Hirth 1993 and 1997), but an updated synthesis of current information is lacking. Seminoff (2004) provided a global status assessment for green turtles, but focused primarily on two index sites in the Pacific (Hawaii and Australia) excluding the numerous small Pacific Island states within Oceania where green turtle nesting also occurs. The 2007 Green Sea Turtle 5-year Status Review (NMFS and FWS 2007) provides a comprehensive summary of green turtle biology, life history, ecology, and population threats but only includes nesting information from four Pacific locations (Hawaii, Guam, Japan, and Australia). The State of the World's Sea Turtles (SWOT)<sup>1</sup> database is another source of information, however, like all datasets, SWOT has its limitations. The database relies on published research and active monitoring projects voluntarily contributing annual nesting data, which in some cases means that known sites will not be represented if projects have not actively provided data to the SWOT effort. Importantly, because there are many known nesting sites without active research or monitoring projects throughout Oceania, the region may be under-represented in SWOT's database.

In an effort to develop a tool for marine turtle resource management, the National Marines Fisheries Service (NMFS) Pacific Islands Regional Office (PIRO) Protected

---

<sup>1</sup> SWOT is a partnership between Conservation International (CI), the IUCN Marine Turtle Specialist Group (MTSG), Duke University's OBIS-SEAMAP, and international collaborators including non-governmental organizations, government agencies, scientists, and conservationists.

Resources Division (PRD) has completed a literature review to better understand green turtle nesting distribution in Oceania. This review also includes applicable sea turtle laws in each country and international arrangements (Appendix I) that exist for sea turtle conservation or management. We recognize that existing legislation and international agreements do not always represent the reality of species protection due to the practical nature of enforcement, compliance, and socio-economic factors that vary widely between nations. While a detailed discussion of legislation's impact on sea turtle conservation is beyond the scope of this document, a spatial summary of legislation is included for reference and generalized conclusions are provided in the discussion.

Marine turtles throughout Oceania have a long history of exploitation and are iconic species of high cultural, traditional, and often spiritual significance to many indigenous groups in the Pacific (Allen 2007; Balazs 1975, 1983a, 1983b, 1995; Campbell 2003; Eckert 1993; Fraizer 2003; Groombridge and Luxmoore 1989; Geermans 1993; Johannes 1978, 1986; Kinan and Dalzel 2004; McCoy 1974, 1982, 1997; Pritchard 1995a, 1995b; Woodrum 2003, 2007, 2010). Conservation is critical not only for persistence of the species, but for preservation of cultural folklore, traditions, and indigenous practices throughout the varied cultures of this region. This synthesis of existing information on Pacific green turtle nesting locations, nesting assemblage characteristics where available, and current legislative protections provides the most current information for the region. It establishes a foundation for future prioritization of monitoring and recovery projects by identifying where current projects occur and where current assessment efforts may be lacking.

## **Methods**

Available information from published documents, gray literature, and in-country government reports of green turtle nesting assemblage locations, sizes (estimates of annual nesting females), and current nesting trends (increasing, declining, stable, or unknown) within each country or territory's exclusive economic zone (EEZ) was compiled. Nesting beach surveys are the most widely implemented monitoring tool used by the global sea turtle community to assess and monitor the status of nesting sea turtle populations (Schroeder and Murphy 1999). For the purposes of this document, the terms monitoring, census, assessment, and survey are used interchangeably and reference any activity employed to gather nesting beach information, such as number of nests deposited or number of nesting females at a location, with no intended implication of duration of the survey, unless noted. Long-term monitoring refers to consecutive-year projects that have employed a standardized monitoring protocol over ten or more years.

This literature review is accompanied by several maps created using ArcGIS™ (version 9.3 software, ESRI, Inc., Redlands, CA) that display the spatial distribution and various attributes of green turtle nesting sites in Oceania. Abundance of nesting females and general trend of the nesting aggregation are described, where possible, based on the best currently available information, and summarized in Table 1. Because the type of information available for nesting green turtle abundance in each country varies significantly, estimates of annual nesting females were binned into the following

categories for the purpose of displaying the information spatially in Figures 1, 1a, and 1b: <10, 10-100, 100-500, 500-1,000, 1,000-5,000, 5,000-10,000, and 10,000-25,000. In cases where virtually no information was available beyond the presence of some green turtle nesting activity, a conservative approach was taken and it was assumed that <10 females nest annually at those sites.

Sea turtle nesting assemblages exhibit natural annual fluctuations in abundance. As such, a conservative approach was taken and a minimum estimated range of annual nesting females was used, unless otherwise noted. Where information was presented in numbers of nests, an estimate of 4.5 nests per individual was applied to determine the number of nesting females likely represented. While this life history trait has not been studied for most of the rookeries in Oceania, our estimate is based on Van Buskirk and Crowder's (1994) reported average for the Heron Island, Australia rookery and an updated estimate for the nesting assemblage at French Frigate Shoals, Hawaii (S. Hargrove, pers. comm.). While the best available nesting information is used in this synthesis, NMFS recognizes that this information represents only one segment (less than 1%) of sea turtle life history (NRC 2010), and therefore population abundance estimates are not inferred from this data. The 155 papers, articles, books, published and unpublished reports, and correspondences used in preparation of this document are included in the References section and linked to the corresponding information within GIS shapefiles.

The generalized legislation reported in this summary is current as of the time of printing, with some possible exceptions where we were unable to confirm the accuracy of the most up to date information with in-country representatives. In these cases, information was adapted from a report compiled by SPREP in 2007; as a result, there may be some instances in which new legislation has been passed since 2007 that is not included in this report. To facilitate comparison of legislation between countries despite nuances of variation, regulations were categorized and presented as an overview within maps, while the text associated with each country provides more detail. Generalized legislative categories include: allowable (permitted/regulated) take; open/closed harvest periods; minimum size limits; protection while on beaches (i.e., nesting females and eggs/nests only); and no legislation specific to green turtles. In many Pacific Island countries, indigenous and subsistence harvest is acknowledged and permitted. We do not discuss the effectiveness or impact of this regulatory exemption and assume that protections are based on indigenous cultural practices such as those described by Woodrum (2003 and 2010), Allen (2007), and other references cited within.

# Synthesis of Existing Data on Green Turtle Nesting and Legislation in Oceania

## ***U.S. States, Territories, and Possessions***

All U.S. states, territories, and possessions are subject to the laws of the United States Federal Government as well as any international agreements to which the U.S. is a signatory. As such, the Endangered Species Act (ESA) of 1973 (16 U.S.C. § 1531 *et seq.*) is the statute under which sea turtles are federally protected in these areas. Six species of marine turtles occurring in the U.S. are listed under the ESA; in the WCPO, five of those species occur in U.S. waters<sup>2</sup> and are under joint jurisdiction of the NOAA Fisheries National Marine Fisheries Service (NOAA Fisheries Service or NMFS) and U.S. Fish and Wildlife Service (USFWS). NOAA Fisheries Service has the lead responsibility for conservation and recovery of sea turtles in the marine environment and USFWS is responsible for sea turtles in the terrestrial environment. Green turtles are listed as *threatened* throughout their range with the exception of breeding colonies in Florida and the Pacific coast of Mexico which are listed as *endangered*. The ESA prohibits unauthorized ‘take’ of listed species which is defined as to “harass, harm, pursue, hunt, shoot, wound, kill, trap, capture, or collect, or to attempt to engage in any such conduct” (16 U.S.C. § 1532(18)). Under the ESA, exceptions to ‘take’ prohibitions are permitted for scientific research or when take is incidental to an otherwise lawful activity, as long as the level of take will not jeopardize the existence of the species in the wild or appreciably reduce the likelihood of recovery in the wild. Both NOAA Fisheries Service and USFWS have dedicated enforcement divisions to handle violations of the ESA. The U.S. is a party to several international agreements related to sea turtles (see Appendix I) including the Inter-American Convention for the Protection and Conservation of Sea Turtles (IAC) and the Convention on International Trade in Endangered Species (CITES) which prohibits all commercial trade of marine turtles and marine turtle derived products. While the U.S. is not a party to the Convention on Migratory Species (CMS), they are a signatory to the Convention on Biological Diversity (CBD). Local state and territory governments may have additional protections in place for sea turtles, as discussed individually below.

## **American Samoa**

The U.S. territory of American Samoa is located east of Western Samoa in the south Pacific and consists of the main island of Tutuila, the Manu`a group (Ofu, Olosega, and Ta`u Islands), Swains Island, and Rose Atoll (Tuato'o-Bartley et al 1993). In American Samoa, sub-adult and adult green turtles occur in low abundance in nearshore waters around Tutuila, Ofu, Olosega, Ta`u and Swains Islands with sporadic, low-level green turtle nesting occurring on Tutuila and Swains Islands (Tagarino et al. 2008; Tagarino and Utzurum 2010). A May 2009 survey at Swains identified a total of 56 locations of pits/possible nests, turtle tracks, and evidence of pig activity (wallows) (Tagarino and

---

<sup>2</sup> Leatherback (*Dermochelys coriacea*), hawksbill (*Eretmochelys imbricata*) loggerhead (*Caretta caretta*), green (*Chelonia mydas*), and olive ridley (*Lepidochelys olivacea*).

Utzurum 2010). However, the primary green turtle nesting location is at Rose Atoll with up to several dozen nests laid annually between October and March (review provided by Balazs 2009). No nesting trend data are available, but anecdotal information suggests major declines in the last 50 years (Tuato'o-Bartley et al 1993, Utzurum 2002). Since 1971, 42 individual nesting green turtles have been flipper tagged on Rose Atoll during various trips (Grant et al. 1997). Of seven post-nesting green turtles satellite-tagged in 1993-95, six migrated nearly directly to Fiji, possibly to feed on Fiji's extensive seagrass beds (Craig et al. 2004). Several surveys cited in a summary of nesting observations at Rose Atoll 1839-1993 (Balazs 2009) documented pits on Sand and Rose Islands (up to 301 in one survey), however, it is unclear how that relates to numbers of individuals because some pits could be from prior nesting seasons.

In addition to protection under the federal ESA, sea turtles in American Samoa are protected by the Fishing and Hunting Regulations for American Samoa which prohibit the import, export, sale, possession, transport, or trade of sea turtles or their parts and take (as defined by the ESA) and carry additional penalties for violations at the local government level. The Department of Marine and Wildlife Resources (DMWR) is the agency with vested authority and responsibility for conservation of protected species and enforcement of protected species regulations in American Samoa.

### **Commonwealth of the Northern Mariana Islands**

The U.S. territory of the Commonwealth of the Northern Mariana Islands (CNMI) consists of 15 volcanic islands in the Marianas archipelago (excluding Guam). The three largest and southernmost islands are Saipan, Rota, and Tinian with the majority of the human population residing on Saipan. In CNMI, green turtle nesting occurs from March through August with some year round nesting documented. It is estimated that possibly fewer than 10 individual turtles nest annually on the islands of Saipan, Tinian and Rota (NMFS and USFWS 1998). Surveys of the northern islands, Alamagan, Pagan, Agrigan, and Asuncion, were sponsored by the Department of Defense and organized by the USFWS from May – June 2010. Turtle nesting activity was only observed on Agrigan, with seven nests documented (C. Eggleston and F. Amidon pers. com.). There were no recorded nesting observations during a survey of Anatahan in 2002 (Ilo and Manglona 2002).

The CNMI Division of Fish and Wildlife (DFW) has monitored nesting activity on Saipan since 1999 and has documented four to eighteen nests laid per year (DFW unpublished annual reports to PIRO). At least five beaches on Saipan have been monitored somewhat consistently over the past five years: Bird Island, Wing, Tank, Lao Lao Bay, and Obyan beaches (Ilo et al. 2005; Kolinski et al. 2001; DFW 2009). Nesting likely occurs on all or most of the accessible beaches on Tinian (Pultz et al. 1999), with six beaches occurring on Navy lands monitored for turtle nesting activity by Navy personnel: Chulu, Lamlam, Babui, Chiget, Dangkulo (Long Beach), and Masalok (Vogt 2009). Eleven beaches on Rota are known to support nesting: Songton, Teteto, Mochong, Sagua (Kokomo), Coral Garden, Okgok, Apanon, and Gaonan (the Cave Beaches), Uyulan, Tatgua, and Latte Stone (Lalayak or I Batko) (Ilo et al. 2005), of which two beaches had confirmed nesting activity in 2009 (Okgok and Tagua).

Intensive monitoring occurred on Saipan at seven beaches from March 4 to August 31, 2009 resulting in 16 green turtle nests documented (DFW 2009). Of major concern, however, is that three of potentially five nesting turtles and three nests were illegally harvested which suggests that poaching remains a significant threat to turtles on Saipan. Rapid assessments at Rota beaches Okgok and Tatgua on July 12, 2009 yielded 13 nests. On Tinian, from July 22-31, 2009, 36 nests at five beaches were documented with evidence of one nesting female illegally harvested (DFW 2009). Additional nesting assessments and dedicated monitoring efforts at Tinian and Rota are needed as these islands may provide viable nesting beaches in CNMI and are likely good candidate index sites for long-term monitoring to assess nesting trends over time. Genetic samples analyzed to date indicate that nesting females in CNMI and Guam are indistinguishable and should be treated as a single management unit (Dutton 2009 unpublished). However, sample sizes are small and additional sampling may reveal other haplotypes. Sufficient information on nesting trend is not available for green turtles in CNMI although anecdotal information from residents suggests that nesting activity has decreased over time, likely as a result of direct harvest, coastal development, and WWII impacts to nesting turtles and their habitats.

In addition to protection under the federal ESA, sea turtles in CNMI are protected by the Fish, Game and Endangered Species Act (PL 2-51). CNMI PL 2-51 establishes a Fish and Wildlife Division and states that the Director of Natural Resources shall determine whether any species shall be designated as threatened or endangered. Green and hawksbill turtles are listed as protected species in the CNMI Hunting Regulations (CNMI DFwb, accessed 2010) prohibiting hunting for these species. The CNMI Department of Land and Natural Resources, Division of Fish and Wildlife is the agency with vested authority and responsibility for the conservation of protected species and enforcement of protected species regulations in CNMI.

## **Guam**

The U.S. territory of Guam is the southernmost island in the Marianas archipelago located in the western Pacific, south of Japan and north of Papua New Guinea. There is regular, low density green turtle nesting on Guam at a number of sites. Nesting activity appears to occur at low levels year round with a more concentrated nesting season apparent from May through August (Pritchard 1995b; NMFS and USFWS 1998a). Documented nesting beaches include: Ritidian National Wildlife Refuge, Haputo, Urunao, Tumon Bay, Cabras Island, the waterfront annex of Naval Base Guam, Spanish Steps, Cocos Island, Acho Bay, Nomña Bay, Jinapsan, and Tarague Beach (DAWR 2004; Grimm and Farley 2008). The nearshore marine environment around Guam has been degraded by impacts from intense combat during WWII, shoreline development, sediment-laden runoff, pollution, and years of poorly treated wastewater effluent. Spanish Steps is at the mouth of Apra Harbor which has been heavily modified, particularly since World War II (USN 2010).

The Guam Department of Agriculture Division of Aquatic and Wildlife Resources (DAWR) initiated a sea turtle program in 1999 with primary objectives to monitor

nesting activity and collect population data. From October 1, 2006 through July 31, 2008, 55 green turtle nests were counted at various beaches during opportunistic surveys throughout Guam (DAWR 2009). Spanish Steps, or Orote point, on U.S. Navy land is considered one of the primary nesting locations on Guam (Grimm and Farley, 2008). Naval Facilities Engineering Command Marianas (NAVFACMAR) monitored nesting beaches at Spanish Steps three times per week from May to July during 2007 and 2008 that resulted in five and 18 green turtle nests, respectively (Grimm and Farley, 2008). Based on this limited information, one to four adult green turtles may nest per season at Spanish Steps; however, sufficient long-term and standardized monitoring information is not available to quantitatively describe the abundance or trend of nesting green turtles at Spanish Steps or for Guam overall. In 2000 and 2007, two post-nesting green turtles were satellite tagged on Guam and traveled to the Philippines and Japan, respectively (NMFS and DAWR unpublished). Currently, nesting activity is documented opportunistically by Haggan-watch, a community-based volunteer network administered by DAWR.

In addition to protection under the federal ESA, sea turtles are protected by the Endangered Species Act of Guam which adopts the same definitions and status designations as the federal ESA and carries additional penalties for violations at the local government level. DAWR is the agency with vested authority and responsibility for the conservation of protected species and enforcement of the ESA of Guam. Other Guam resource agencies, such as the Bureau of Statistics and Plans (BSP), also have specific mandates in relation to sea turtle conservation. The BSP administers the Guam Coastal Management Plan (GCMP) through the Coastal Zone Management Act of 1972 (Guam Public Law 92-583 and Public Law 94-370). The GCMP guides the use, protection, and development of land and ocean resources within Guam's coastal zone, which includes all non-Federal property and all submerged lands and waters out to 3 nm (5.6 km) from the shoreline.

## **Hawaii**

The State of Hawaii is an archipelago in the central Pacific Ocean containing hundreds of volcanic islands, separated into two groups: eight large southeastern Main Hawaiian Islands (MHI; seven of which are inhabited), and numerous uninhabited Northwestern Hawaiian Islands (NWHI; designated the Papahānaumokuākea Marine National Monument by Presidential proclamation in June 2006). Green turtles nesting and foraging within the Hawaiian Archipelago are likely comprised of one genetic stock, and may be considered a discreet management unit separate from other Pacific stocks (Dutton et al. 2008). Nesting occurs between May and August, and the primary nesting location at French Frigate Shoals (FFS) in the NWHI supports over 90% of documented green turtle nesting in Hawaii (Balazs 1976, 1980). Minor nesting also occurs at other atolls and islands in the NWHI<sup>3</sup> and on Kauai, Oahu, Molokai, Lanai, and Maui within the MHI (PIFSC unpublished). Within FFS, over 50% of all nesting occurs on East Island (Balazs 1976; Niethammer et al. 1997, Balazs and Chaloupka 2004), where nesting surveys have

---

<sup>3</sup> Known nesting occurs at Laysan, Lisianski, Pearl and Hermes, and Midway. Four infertile nests were laid at Kure in 2009. No information exists about nesting activity at Nihoa and Necker.

been conducted annually at this index site since 1973 via a collaborative arrangement between NMFS Pacific Islands Fisheries Science Center (PIFSC) and USFWS.

The Hawaiian green turtle population was subjected to extensive human exploitation in the form of turtle and egg harvest at foraging and nesting grounds from the mid-1800's until the early 1960's, and nesting habitat destruction as a result of development (Balazs 1975a, 1976; Niethammer et al. 1997; Balazs and Chaloupka 2004).<sup>4</sup> Since enactment of State and federal ESA protections in 1974 and 1978, respectively, the nesting population at FFS has exhibited high annual variability in nesting female abundance, and a consistent upward trend over the past thirty years with an estimated annual growth rate of 5.7% (Chaloupka et al. 2008). The largest number of nesting females observed during a field season at East Island occurred in 2008 with 580 females identified during the six week sampling period (PIFSC and FWS unpublished).

In addition to protection under the federal ESA and international agreements and conventions, sea turtles in Hawaii are protected by the Hawaii Revised Statutes, Chapter 195D (Hawaii State Legislature, accessed 9/10/2010) and Hawaii Administrative Rules, 13-124 (Hawaii Administrative Rules, accessed 9/10/2010) which adopt the same definitions, status designations, and prohibitions as the federal ESA and carry additional penalties for violations at the State government level. The Hawaii Department of Land and Natural Resources (DLNR) Division of Aquatic Resources (DAR) is the state agency responsible for the conservation and management of protected species in Hawaii. The Division of Conservation and Resources Enforcement (DOCARE) is the agency with enforcement authority at the state level in matters involving violations of Hawaii's protected species regulations.

### **Pacific Remote Island Areas**

The Pacific Remote Island Areas (PRIAs) are U.S. possessions that are widely spread throughout the Pacific and include Wake, Johnston and Palmyra Atolls, Kingman Reef, and Jarvis, Howland, and Baker Islands. Following a 28 day assessment in 1983 it was concluded that green turtles do not nest at Johnston Atoll, but occur foraging within the atoll (Balazs and Forsyth 1986). Low-level nesting was observed at Palmyra in 1987 and along the west coast of Jarvis Island in the 1930s (NMFS & FWS 1998) but no recent surveys have been conducted. Both Jarvis and Palmyra are geographically part of the Line Islands chain of coral atolls and islands in the central Pacific and are uninhabited remote National Wildlife Refuges administered by the USFWS. Jarvis is visited infrequently by refuge staff for one to two days at a time every two years. There is a research station on Cooper Island at Palmyra Atoll operated by The Nature Conservancy (TNC) that houses a small maintenance staff year-round and various research groups for shorter time periods. Anecdotally, no evidence of sea turtle nesting has been observed at Palmyra in recent years (USFWS, pers. comm.). In 2007, an in-water sea turtle research project was initiated at Palmyra by the American Museum of Natural History and

---

<sup>4</sup> Green turtle and egg harvest at FFS officially ended in 1964 when it was permanently placed under the jurisdiction of the USFWS, but the last documented harvest event was in 1959 when a commercial fishing boat harvested a minimum of 25% of nesting females present that season (Balazs 1975a).



Columbia University. While nesting beach monitoring is not a focus of the project, any nesting activities will be documented by either the project or by TNC staff that currently reside at the Atoll.

The PRIAs do not support resident human populations and do not have local governments. Therefore, all sea turtle species that occur in the PRIAs are protected by the federal ESA as described previously.

## ***International – States with Compacts of Free Association<sup>5</sup>***

### **Federated States of Micronesia**

The Federated States of Micronesia (FSM) consist of 607 islands in the western Pacific that are divided into four states: Yap, Chuuk, Kosrae, and Pohnpei. Green turtle nesting in FSM peaks between April and August, with some evidence of year round nesting (J. Cruce pers. comm.), in all states except Kosrae where nesting has not been documented. FSM has laws and regulations that prohibit the harvest of all sea turtles while they are on shore and of all eggs, establishes minimum size limits for hawksbills (27 inches CCL) and greens (34 inches CCL), and establishes closed seasons from June 1 – August 31 and December 1 – January 31, during which time no sea turtle of any size shall be taken or killed ([www.fsmlaw.org](http://www.fsmlaw.org), accessed 9/10/2010). However, these regulations are only applicable within FSM waters outside the 12 mile-state territorial waters zone. FSM is not a signatory of CITES. Each state has additional individual regulations as described below.

Yap State, FSM is comprised of approximately 134 islands and 11 atolls of which 22 are inhabited by Yapese people, many of whom continue to practice cultural traditions. Turtles are an integral part of many aspects of Yapese life. Green turtles are the most common species nesting in Yap. During a four and a half month field season on Olimarao Atoll in 1990, 27 adult females were tagged (Smith et al. 1991). A field season on Elato Atoll in 1992 yielded 36 tagged adult females (Kolinski 1993). A total of 70 nesting green turtles were tagged in Ngulu Atoll on the islands of Lathow and Meseran from May through July in 1992 and another 75 nesting green turtles were tagged the following year from April through July of 1993 (Kolinski 1993). In addition, two tagging efforts were carried out at Elato Atoll where 36 nesting green turtles were observed from July through September of 1992 and 41 nesters from May through August of 1993 (Kolinski 1993). Ulithi Atoll, located approximately 185 kilometers (km) northeast of Yap Proper, is home to several “Turtle Islands” which are identified as significant green turtle nesting sites by local people including the island trio of Loosiep, Bulbul and Yeew and duo of Gielop and Iar (Cruce 2006). These islands may be among the largest green turtle rookeries in Micronesia (Kolinski 1992) and are the focus of current monitoring efforts in

---

<sup>5</sup> “Freely Associated States”, the Republic of Palau, Federated States of Micronesia and Republic of the Marshall Islands, were formerly governed by the U.S. as part of the United Nations Trust Territory of the Pacific Islands but have since become sovereign states, are still eligible to receive funds from U.S. Federal agencies, and maintain close ties with the United States.

Yap. Turtles nesting on or mating near these islands have traditionally been hunted for their meat and eggs (Lessa 1983).

In 1991, 417 nesting green turtles were tagged on Gielop between May and August (Kolinski 1992). This study site was revisited subsequently when a turtle tagging project was carried out on the islands of Gielop and Iar from June 9 to August 24, 2005. Islands were monitored a total of 59 nights on Gielop and 25 nights on Iar with a total of 310 nesting green turtles tagged (186 from Gielop and 124 from Iar) and one nesting hawksbill turtle on Iar (Cruce 2006). During subsequent monitoring seasons on Gielop in 2006 and 2007, 329 and 250 nesting green turtles were tagged, respectively, between June and August each year (Cruce 2007).

In 2008, a research ban was instituted by local chiefs at Gielop so monitoring efforts switched to nearby Loosiep Island where 66 nesting green turtles were tagged between April 22 and July 18 with some evidence of predation documented (Cruce 2008). Research resumed at Gielop in 2009, with a total of 553 green turtles recorded, including eight non-nesting female green turtles and one male turtle tagged and assessed. Tag returns were documented from turtles originally tagged on Gielop in 2005 and 2006. At Loosiep Island, 118 nesting females were tagged between April 13 and August 2 with continued evidence of nest predation by exotic varanids and pigs (17 of 20 staked study nests) suggesting that management measures are needed on this island (Cruce 2009).

Genetic samples have also been collected during monitoring efforts, of which 462 have been analyzed to date. Results suggest nesting green turtles in Yap are comprised of one genetic stock distinguishable from other Pacific nesting populations, although additional samples and analysis are needed to provide increased resolution (Dutton 2009 unpublished). Of seven post-nesting green turtles satellite tracked from Gielop during 2005-2006, five migrated to the Philippines and one to Malaysia while another turtle's transmitter ceased sending signals while still in the FSM EEZ (Kolinski et al. 2007 draft manuscript). An additional seven post-nesting greens were tracked from Gielop in 2007; four turtles migrated to the Ryukyu Islands, Japan and three to the Philippines (PIRO and PIFSC unpublished data).

Yap State Code prohibits the commercial sale of sea turtle meat and eggs (Yap State Code, accessed 9/10/2010). Traditionally, nesting green turtles throughout Ulithi Atoll have been managed and conserved by the imposition of cultural limitations on take for consumption, put in place by leaders of the chief island, Mogmog (Lessa 1983). In recent years, it appears turtle take has increased due to the degradation of traditional practices although the number of turtles taken annually within Ulithi Atoll has not been assessed or quantified (Cruce 2009).

In Chuuk State, FSM, nesting sites include Fanang Island, East Fayu Island, and Murilo Atoll. According to a report from 1993, six to seven green turtles nested each night from February to June on East Fayu (Pritchard 1995b). One to three turtles per night were also reported to nest on Fanang Island as well as a few per night at Murilo Atoll (Pritchard 1995b). Chuuk State Code was still in draft form at the time of drafting of this report.

In Pohnpei State, FSM, green turtle nesting has been recorded at Oroluk Island, Oroluk Atoll. Oroluk is an atoll west of Pohnpei Island and has over 30 sandy islets and sandbanks. Only Oroluk Island is inhabited with fewer than 20 residents. Pritchard (1977) noted that Oroluk Atoll was apparently the only nesting ground of importance for the green turtle in Ponape District (Pohnpei State) and he estimated 9 to 15 nests per night at Oroluk with up to 20 nests on a good night. Pritchard also reported a nesting pattern with two peaks, December to January and June to July. At least some nesting is reported by inhabitants all year round (Edson and Curren 1987). Surveys from June through July of 1985 resulted in an average of 2.3 nests per month and May through August of 1986 averaged 3.4 nests per month, significantly lower abundance than the 9 to 15 nests per night reported by Pritchard in the 70s (Edson and Curren 1987). During a one day survey in November, 1990, no evidence of nesting was seen on Oroluk, however, Typhoon Owen had passed just north of the island eight days earlier and caused considerable damage to the island and reefs (Naughton, 2001). During Naughton's 1990 expedition, an individual on the island stated that between five and eight turtles nest or attempt to nest on Oroluk Island every month, except June and July when they are "too numerous to count." He reported that island residents take every turtle they encounter. In the 1990s, nesting activity still occurred on Oroluk although at a reduced level from that reported in the 1970s. According to Naughton (2001), there is little question that Oroluk Atoll is critically important to green turtles in the Caroline Islands, (Eastern range of FSM) and is probably the most important site for the species in eastern FSM.

Regarding other sites in Pohnpei, 74% of people surveyed by Buden and Edwards (2001) on Pohnpei Island indicated they had no knowledge of nesting activities of turtles on Pohnpei and its lagoon islands. Four people contributed unsolicited comments on nesting on Ant Atoll, and three described incidents of nesting on three different lagoon islands in Kitti (Buden and Edward 2001) indicating the possibility of very low-level nesting at a few sites in addition to Oroluk Atoll.

According to Pohnpei State laws, there is a minimum size limit for greens (34 inches carapace length) and hawksbills (27 inches carapace length) and closed harvest seasons June 1 to August 31 and December 1 to January 31, with nesting turtles protected and egg collecting prohibited at all times (SPREP 2007; Buden and Edwards 2001).

Although no nesting has been reported in Kosrae, state code regulates the take of turtles in water with a minimum size limit for all species of 27 inches carapace length, closed seasons June 1 to August 31 and December 1 to January 31, and prohibition of egg collecting and killing turtles while onshore at all times.

It is estimated that between 500 and 1000 green turtles nest annually in FSM, however, estimates are based on available data from the few nesting sites that have been monitored and sampled, whereas green turtles may nest at many more sites throughout Micronesia undocumented. As such, it is likely that we have underestimated nesting activity in this under-monitored region. The unknown impact of long-term and unregulated harvest of nesting females and their eggs in FSM further supports the need for more consistent

monitoring and additional monitoring sites to establish the current status of the nesting green turtle population within FSM.

## **Republic of the Marshall Islands**

The Republic of the Marshall Islands (RMI) is made up of 29 atolls and five islands with a total land area of approximately 70 square miles, and a total lagoon area of about 4,500 square miles. Atolls and low coral islands are aligned in two roughly parallel northwest-southeast chains: the northeastern Ratak Chain and the southwestern Ralik Chain. Green turtles are most common in the RMI with hawksbill turtles considered rare or scarce (NMFS and FWS 1998). Atolls most recognized as significant green turtle nesting areas include Bikar, Erikub and the island of Jemo. Additional minor nesting sites include the atolls of Bokak, Ailinginae, Rongerik, Bikini, Wotje, and Taka (McCoy 2004). First described by Tobin (1952 in McCoy 2004), northern RMI atolls are well known traditionally as “game reserves” due to the presence of nesting turtles and seabirds (this refers to Bikar, Bokak, and Taka atolls, the island of Jemo, and certain islands in Erikub atoll). Nesting occurs from May through November, peaking mid-June to mid-September. Lagoons throughout Marshall Islands atolls provide significant areas of potential shallow water foraging habitat for sea turtles (Eckert 1993), but in general, sea turtle nesting and foraging activity are more common in inverse proportion to proximity or density of human habitations and activities in the RMI (McCoy 2004).

Bikar Atoll likely supports the largest green turtle nesting assemblage in the RMI. Based on Hendrickson’s observations in 1972 (cited in McCoy 2004), approximately 950 nests were laid, or 237 females may have nested annually at Bikar. At the time, Hendrickson concluded that Bikar represented one of the major breeding groups of sea turtles in the then-Trust Territory of the Marshall Islands (McCoy 2004). NMFS and FWS (1998) estimated a mean annual total of approximately 100-500 nesting females at Bikar Atoll based on an 11 night survey where 48 turtles were tagged in 1992. During the same expedition, eight turtles were tagged in one night on Jemo and a one-time survey of Erikub Islet at Erikub Atoll revealed “. . . many nesting excavations, some well within the interior of the islet. So numerous were these excavations that no attempts were made to count them” (Puleloa and Kilma 1992). Also at Erikub, two pits were observed on Aradojairek Islet and 48 pits on Aradojairen Islet, although it was apparent that some of these were from previous seasons.

Five post-nesting green turtles were satellite tagged on Loj Islet, Erikub Atoll in 2007-2008. One turtle migrated to Bikini Atoll, RMI, one to Tarawa Atoll in Kiribati, one to the Philippines passing through CNMI, and two turtles circled for long periods of time in the open ocean with one having final transmissions in the FSM EEZ and the other in RMI (PIRO and PIFSC unpublished). Between 2005 and 2007 a project supported by NMFS PIRO was implemented to collect genetic samples. Results from the analysis of 125 samples suggest that green turtles nesting in RMI are comprised of one genetic stock distinguishable from other Pacific nesting populations in Palau, Yap and the Mariana Islands (Dutton 2009 unpublished). Again, additional samples from nesting sites throughout RMI are needed to provide greater resolution for a regional stock analysis.

Turtles in the RMI have long been known as a food source and have played an important cultural role in the lives of inhabitants. In 2009 the Marshall Islands Marine Resources Authority began to implement an outreach and education program with support from NMFS PIRO, which may represent the first concerted effort to conserve this cultural resource in the RMI. The level of exploitation of turtles is unknown, and there are no current data available on status of turtle stocks in the RMI (McCoy 2004), although based on the available information an estimated 100-500 green turtles may nest in the RMI annually (NMFS and FWS 1998). While there does not appear to be enough data to conclude if trends are increasing or decreasing, anecdotal information from local people suggests that the number of nesters has decreased over time, possibly by as much as 50 percent in the last 10 years (McCoy 2004).

The harvest of sea turtles in the RMI is regulated by the Marine Resources Act (RMI 1997) which sets minimum size limits for greens (34 inches carapace length) and hawksbills (27 inches carapace length) and closed seasons from June 1 to August 31 and December 1 to January 31. Egg collecting and take of turtles while they are onshore is prohibited at all times. The Marshall Islands Marine Resources Authority is the entity with the responsibility of managing marine resources in the RMI. RMI is not a participating party to CITES.

## **Republic of Palau**

Palau is an island nation made up of four populated islands and several hundred smaller islands and atolls organized into 16 states, 800 km east of the Philippines. Green turtle nesting has been documented at Helen Reef in Hatohobei State and Merir Island in Sonsorol State with additional low level nesting in Ngarchelong, Kayangel, and Melekeok States. Based on 42 genetic samples analyzed to date in comparison with samples from nearby rookeries, green turtles nesting at monitored sites in Palau appear to be one genetic stock distinguishable from other Pacific nesting populations, although additional samples are needed to provide increased resolution (Dutton 2009 unpublished). Nesting summaries for each State are as follows:

Hatohobei State: During a study at Helen Reef from April 19 through December 8, 2005, 301 green turtle nests were counted, 47 individual nesting turtles were flipper tagged, and nesting turtles emerged almost every night between April and August 2005 (Barr 2006). On April 22, 2008 a female green originally tagged on Helen Island, Hatohobei State, on September 5, 2006 was recovered near Goulburn Island, Northern Territory, Australia after being speared with a traditional harpoon (Palau BMR 2008). Reduced abundance of green sea turtles at Helen Reef has been noted by the Tobian community (Emilio et al. 2002 in Barr 2006).

Sonsorol State: From November 2004 through September 2005, 331 green turtle nests were documented during daily surveys and 36 individual turtles were tagged during night surveys at Merir Island (Palau BMR 2005). Five green turtle nests were documented during surveys conducted on April 17 and 25, 2005 at Pulo Ana Island (Palau BMR 2005). During daily monitoring from November 2007 to August 2008, 739 green turtle nests and 382 non-nesting emergences were documented with peak nesting observed in

May (Palau BMR 2008). A green turtle tagged on June 7, 2007 by conservation officers on Merir Island, Sonsorol State was recaptured in a set net near the village of Yomitami, Okinawa, Japan on October 15, 2007. The turtle was retagged and released (Palau BMR 2008).

Ngarchelong State: Between March and August of 2008, four surveys found 11 green turtle nests along 2.4 km of the island of Ngerechur, nine of which had been destroyed by wild pigs (Palau BMR 2005). On July 22 and 23, 2008 turtle nesting surveys were conducted along a 1.5 km beach on Ngerechur Island, just off of Ngarchelong state where one green turtle nest was documented (Palau BMR 2008).

Kayangel (Ngcheangel) State: Kayangel is an atoll with four islets on its east side. Kayangel Island, with a land area of 1.12 km<sup>2</sup> and a perimeter of 6 km, is the largest island in the atoll. Ngeriungs is just south of Kayangel and is the second largest island of the atoll with a land area of .32 km<sup>2</sup> and a perimeter of 3 km. Between April 28 and October 10, 2005, two green turtle nests were documented during occasional nesting beach surveys on Kayangel Atoll (one on Kayangel and one on Ngeriungs) (Palau BMR 2005). Green turtle nesting also occurs at Ngeruangel Islet, Ngeruangel Atoll, 10 km northwest of Kayangel. Between June 22, 2005 and October 10, 2005, five green turtle nests were documented as a result of three surveys (Palau BMR 2005).

Melekeok State: Melekeok is a town on the east coast of Babeldaob Island with a beach area of 4.43 km<sup>2</sup>. On, November 11, 2005, five sites along the beach were surveyed with no turtle nests documented. Interviews with several residents indicated turtle nesting in the area with a maximum of five green turtle nests in a year (Palau BMR 2005).

Palau domestic fishing laws specify minimum size limits for green turtles (34 inches carapace length) and hawksbills (27 inches carapace length) and closed seasons from June 1 to August 31, and December 1 to January 31 (SPC and BMR Palau, 2007). Taking of eggs or female turtles while onshore is prohibited at all times. Palau is a participating party to CITES.

## ***International***

### **Australia**

Green turtles nest at numerous sites along the east, north and west coasts of Australia as well as many islands and islets offshore and on the Great Barrier Reef (GBR) and in the Coral Sea. To remain within the scope of this document, only those sites along the eastern coast, on the GBR and in the Coringa-Herald National Nature Reserve (CHNNR) in the Coral Sea will be discussed. Nesting occurs in these areas between October and March (Limpus 2009). The Northern Great Barrier Reef (NGBR) includes the largest nesting concentration of green turtles in the world (Chaloupka et al. 2007), with 90% of nesting activity in the area occurring on Raine Island, Moulter Cay, and No. 7 and No. 8 Sandbanks (Limpus 2009). Minor breeding aggregations also occur on the Murray Islands, Bramble Cay, and other outer barrier islands of the NGBR, most inner shelf cays,

and mainland beaches north of Cape Grenville and along the Torres Strait (Limpus 2009). Raine Island is the primary index beach for the NGBR stock, but a total tagging census has not been attempted and there has been limited annual monitoring of the nesting aggregation at NGBR and Torres Strait rookeries due to size of the nesting assemblage and logistical challenges associated with site remoteness (Limpus 2009). This region experiences significant inter-annual fluctuation in nesting female abundance, ranging at certain sites from a few dozen to over 10,000 annual nesting females, driven primarily by the El Niño Southern Oscillation (Limpus et al. 2003; Chaloupka et al. 2007; Limpus 2009). Moulter Cay has nightly nesting activity and average density that is strongly correlated with the activity and density at Raine Island. High density nesting (in excess of 10,000 females annually) was reported in the past in the mid 1950's and 60's (Limpus 2003). It is also expected that during a high density season several thousand additional females nest at No. 7 and No. 8 Sandbanks, the Murray Islands, Bramble Cay and other smaller nesting sites in the NGBR and Torres Strait (Limpus 2009). Although Limpus (2009) does not estimate size of the adult female population occurring at Raine Island, he approximates that 41,000 female green turtles may breed annually during a typical high-density nesting season. Seminoff (2004) and NMFS and USFWS (2007) estimate 18,000 and 25,000 annual nesting females at Raine Island respectively. Based on this available information, a reasonable conservative estimate of the annual mean number of nesters in the NGBR is 10,000-25,000 females.

Chaloupka et al. (2007) identified a nonlinear nesting trend, increasing from the mid-1970s and leveling off by the mid-1990s. Lack of a continued increasing trend at Raine Island may be due to a number of factors including: increasing sea surface temperature (Chaloupka and Limpus 2001; Limpus et al. 2003); decreasing reproductive output as the stock approaches carrying capacity (Troëng & Chaloupka 2007); over-harvest in northern Australian and New Guinean waters (Limpus et al. 2003; Limpus 2009); and hydrology or rising groundwater that floods egg chambers (Limpus et al. 2003). Therefore there is concern regarding long-term stability of the NGBR nesting assemblage given a significant decline in breeding success (low hatchling production and recruitment) over the last three decades at Raine Island (Limpus et al. 2003; Limpus 2009). Additionally, there has been a significant downward trend in mean curved carapace length (CCL) of nesting females at Raine and Moulter Cay over 26 breeding seasons, 1976-2001 (Limpus et al. 2003). This decrease in carapace size has occurred in conjunction with a progressive increase in remigration interval (Limpus 2009), and while long-term monitoring for abundance of annual nesters has not provided a clear indication of the stability of this stock, changes in CCL and remigration interval are consistent with a group that could be in early stages of decline as a result of excessive loss of adult females (Limpus et al. 2003; Limpus 2009).

In the southern Great Barrier Reef (SGBR), major green turtle breeding areas include the islands of the Capricorn Bunker Group: Northwest, Wreck, Hoskyn, Tryon, Heron, Lady Musgrave, Masthead, Erskine, Fairfax, North Reef, and Wilson Islands. Minor breeding aggregations occur at Bushy Island, the Percy Islands, Bell Cay, Lady Elliot Island, mainland beaches from Bustard Head to Bundaberg, and the northern part of Fraser Island. Greater than 90 percent of all SGBR nesting occurs within protected habitats of

National Parks and Conservation Parks (Limpus 2009). Size of the annual breeding assemblage has been monitored at several rookeries for varying periods since 1964 and there exists a wealth of information for this stock (summarized in Limpus 2009). Heron Island is the SGBR index nesting beach and has exhibited a stable fluctuation (i.e., no significant upward or downward trend) in annual nesting activity for almost four decades, 1967-2004 (Limpus 2009). However, there has been significant long-term reduction in mean size of breeding females within this stock over 26 breeding seasons that may be indicative of over-harvest of adult females (Limpus 2009). Based on mid-season nightly track counts, the SGBR is estimated to support 5,000-10,000 nesting green turtles annually (Limpus et al. 1984 and Limpus 1985, in Limpus 2009).

The Coringa-Herald National Nature Reserve (CHNNR) is located 440 km east of Queensland, Australia and is comprised of three pairs of islets: Herald Cays (NE & SW), Coringa Islet (SW & Chilcott), and Magdelaine Cays (NW & SE). Nesting takes place at NE Herald Cay, SW Herald Cay, Chilcott Islet and SW Coringa Islet. The reserve is afforded some protection by virtue of its remoteness and lack of introduced predators (Harvey et al. 2005). Nesting activity at CHNNR extends from late October until approximately mid April, peak nesting occurring from late November through February. With the exception of the 1992/93 nesting season, NE Herald Cay was monitored for 13 years from 1991/92 through 2003/04 with surveys that ranged from four to 33 nights per season. Additionally, SW Herald Cay, Chilcott Islet and SW Coringa Islet were monitored sporadically during this 13 year timeframe. A total of 6,193 female turtles were recorded nesting at all four islets, 4,924 of which nested at NE Herald Cay. Yearly nesting abundance ranged from 12 females (2000/01 season) to 1,445 females (1999/00 season) (Harvey et al. 2005). During the 13 year monitoring period 3,141 turtles were tagged, 2,267 of which were tagged at NE Herald Cay (Harvey et al. 2005). Moritz et al. (2002) report 1,095 green turtles tagged during the 1999/00 season in the CHNNR. In the same season, Harvey et al. (2005) recorded 1,715 total nesting turtles during 33 survey nights of which 922 individual turtles were tagged at three islets: NE & SW Herald Cays and SW Coringa Islet. Insufficient data are available to discern an overall nesting trend, however curved carapace length of nesting females has declined significantly over time which may be the result of harvest pressure or other sources of adult mortality, potentially from the Torres Strait/Papua New Guinea region (Harvey et al. 2005).

Marine turtles in Australia are protected under the Environment Protection and Biodiversity Conservation Act of 1999 (EPBC Act), which implements several international agreements/conventions to which Australia is a signatory. The EPBC Act lists all marine turtles in Australia as 'threatened' species, and provides several mechanisms to address declines in population numbers of listed species that include: recovery plans, threat abatement plans, wildlife conservation plans, conservation agreements and conservation orders. Australia is a participant in the Convention for the Protection of the World Cultural and Natural Heritage (World Heritage Convention), as well as CMS and CITES, both of which list sea turtles under Appendix I (species threatened with extinction). Traditional Owners, as recognized under the Australian Government's Native Title Act of 1993, are able to assert their rights to gain customary



authority for shared resources such as marine turtles which includes traditional hunting rights. Within the Great Barrier Reef, Traditional Use of Marine Resources Agreements can be accredited under Marine Park zoning plans, and Indigenous Land Use Agreements are also being used more frequently. Indigenous groups in Australia have made progress working with governments, wildlife managers and scientists to develop co-operative frameworks of shared expertise and decision-making that incorporates their cultural, social, and political imperatives (K. Dobbs pers. comm.). These initiatives are intended to give indigenous groups greater authority and responsibility, and hence a sense of ownership, commitment, and compliance with regulations as opposed to government-imposed initiatives to sustainably manage turtle resources.

## **Cook Islands**

The Cook Islands consist of fifteen islands spread over 2.2 million square kilometers of ocean, divided into two distinct groups: the Southern Cook Islands which were formed by volcanic activity, and the Northern Cook Islands which are older and consist of six atolls.

In the northern Cook Islands, green turtles nest at Penrhyn (Tongareva), Rakahanga, Pukapuka, and Manihiki Atolls (Balazs 1995). Woodrum-Rudrud (2010) additionally lists green turtle nesting activity at Suwarrow atoll and Nassau island, although additional information regarding number of nesting females is not included for these sites. At Pukapuka, green turtles nest on one of the uninhabited islets and there is some directed harvest of turtles and eggs (Balazs 1995). Reports from the 1960s and 1970s indicate the presence of green turtle nesting activity at other locations but no further details on nesting female abundance or trends are available (Balazs 1995). In the southern Cook Islands, green turtles nest primarily at Palmerston Atoll, which hosts the majority of green turtle nesting within the Cook Islands. According to a review provided by Balazs (1995), reports from the 1960s and 1970s refer to Palmerston as an important nesting location for green turtles in the Pacific, although no indications of numbers of nesting females were provided. From 1972 to 1977 a decline in the number of nesting turtles was observed by inhabitants (Balazs 1995). Annual nesting numbers declined from 30 or 40 to fewer than 10 in under ten years (Helfrich 1974). Additional sites in the southern Cook Islands identified by Woodrum-Rudrud (2010) include Mangaia, Atiu, Mauke, and Roratonga islands, although no further information on nesting abundance is available.

In November 2000, the Center for Cetacean Research and Conservation, a local NGO in the Cook Islands, conducted a consecutive 25 night nesting beach survey on Cook's Islet of Palmerston Atoll (CCRC 2004). Only seven turtles were recorded nesting at Cook's Islet during this survey, and few turtles crawled at the other islets, indicating much lower level nesting than previously reported (anecdotally) at this site. These results were corroborated by Palmerston fishermen, all of whom commented on the paucity of turtles that season (CCRC 2004).

In the Cook Islands, the Marine Resources Act of 1989 provides for the protection and management of fishery resources, the definition of which includes marine turtles. Specific regulations regarding marine turtle harvest were unavailable at the time of

printing although there is full protection of marine turtle eggs (Pulea 1992). The Cook Islands is a participating party to CITES.

## **Fiji**

Fiji consists of an archipelago of more than 300 islands, of which 110 are permanently inhabited, and more than 500 islets located in the South Pacific. The last remaining nesting sites for green turtles in Fiji are small, isolated islands and sand isles north of Taveuni including Nanuku Levu and Nukumbalati Islands within the Hemskercq and Ringgold reef systems. In 1970, eight nests were observed and in January 1980, 16 nests were observed at Nanuku Levu and Nukumbalati (Guinea 1993). As of 1996, the Fisheries Division estimated 30 to 40 nesting green turtles in Fiji (Weaver 1996) with a more recent estimate of 50 to 75 (Batibasaga et al. 2006). A commercial ban on sea turtle harvest was instituted in 1997 (Batibasaga 2002). However, green turtles in Fiji are regularly harvested for consumption and harvest continues to play a significant role in the subsistence economy of many Fijian communities despite the moratorium [May 1997 to December 2000, and February 2004 to December 2008, recently extended through 2019] (Guinea 1993; Laveti and MacKay 2009). There are no long term studies in Fiji to provide information on sea turtle nesting trends but evidence suggests a decline in nesting green turtles due mainly to overharvest (Batibasaga et al. 2006).

Sea turtles and their eggs are managed under Fisheries Regulations in Fiji. The Fisheries Act, as amended in 1979 and 1991, outlines gear requirements when spearing a turtle and also states that “No person shall at any time dig up, use, take, sell, offer or expose for sale, or destroy turtle eggs of any species or in any way molest, take, sell, offer or expose for sale, or kill any turtle the shell of which is less than 455 mm [18 inches] in length. No person during the months of January, February, November or December in any year shall in any way molest, take, sell, offer or expose for sale, or kill any turtle of any size”. The possession, sale or export of any turtle shell less than 18 inches in length and the export of turtle flesh and turtle shell unless it is worked into jewelry or otherwise processed into a form approved by the Permanent Secretary for Primary Industries and Cooperatives are all prohibited acts (Government of Fiji 1992). A National Moratorium prohibiting the killing, harming or molesting of any marine turtles including their meat, eggs or shell was first enforced in 2004 by the Fijian Government. It was recently extended for a further ten years by the Fijian Cabinet until 2019. Indigenous Fijians are still able to legally harvest marine turtles if they obtain prior approval from the Fisheries Department (South Pacific Projects, accessed 9/10/2010). There is some disagreement, however, regarding the effectiveness of Fiji’s moratorium mostly due to lack of compliance and enforcement (Laveti and Mackay 2009). Fiji is a participating party to CITES.

## **French Polynesia**

French Polynesia consists of 130 islands and atolls spread over a large geographic area in the central south Pacific. In western and central French Polynesia, green turtles have historically been observed nesting at Tupai Atoll, Maupiti, Bellinghausen Atoll, Manihi Atoll, Tetiaroa Atoll, Bora Bora, Mopelia Atoll, and Scilly Atoll. Green turtle nesting is concentrated at Scilly Atoll in the Leeward Islands. Observations in the late 1970s, early 1980s, and early 1990s suggested 300 to 400 nesting females occurred there annually

between November and March (Balazs et al. 1995; Lebeau 1985). These observations in conjunction with information from local residents indicate a likely decline in nesting numbers between the 1950s and early 1970s, although numbers may have stabilized between 1972 and 1991 (SPC 1979b; Balazs et al. 1995; Pritchard 1995a). Nesting females and adult males flipper tagged and/or satellite tagged at Scilly Atoll have been recovered in Tonga, New Caledonia, Vanuatu, the Cook Islands, and Fiji (Balazs et al. 1995; Craig et al. 2004); this tag return information reveals some of the longest range migrations recorded for green turtles (SPC 1979b, Craig et al. 2004).

Nesting occurred on Manihi Atoll in 1971 (Hirth 1971, cited in Pritchard 1995a) but recent information is not available. A survey in 1991 at Bellinghausen Atoll resulted in 33 tagged green turtles (SPREP TRENDS 2009 Annual Report). Sporadic nesting surveys at Tetiaroa Atoll have been conducted since 2004 although 2008-2009 was the first nesting season with an organized, sustained survey effort which revealed 81 crawls and 33 nests (Te Mana O Te Moana 2009). Low level nesting has also been observed at Tikehau Atoll (Te mana Tea 2007), and four to 18 turtles nest each year on Bora Bora (Chelonia Polynesia, pers. comm.). Since 2000, adult green turtles occurring on Maupiti Island and Tupai Atoll (Leeward Islands) have been harvested<sup>6</sup> (Chelonia Polynesia, pers. comm.). During the 2007 nesting season, only one crawl was recorded at Tupai.

In eastern French Polynesia, green turtles have been documented historically nesting at Pukapuka Island (a different island from Pukapuka coral atoll in the northern Cook Islands). The most recent information is from 1938 and does not provide an estimate of annual nesting females, although it is noted that turtles and eggs were regularly taken for consumption and residents were already beginning to observe turtles “dying out” (Beaglehole and Beaglehole 1938, cited in Pritchard 1995a).

French Polynesia is an overseas territory of France and sea turtles have been completely protected since 1990 by the Polynesian government (DELIBERATION No. 90-83 AT du 13 Juillet 1990 relative à la protection des tortues marines en Polynésie Française). Prior to this date, traditional harvest with seasonal and size restrictions was permitted. Under the revised statutes, turtles are fully protected and it is strictly forbidden to harm, own or hunt sea turtles or engage in commerce of any kind pertaining to the sale of shell, meat and eggs. Scilly Atoll has been protected as a marine reserve for sea turtles since 1971 by the local government (Vu l'arrêté No. 2559 DOM du 28 Juillet 1971 portant classement du lagon de l'île Manuae ou Scilly). French Polynesia is not a participating party to CITES (although France is a participating party).

## Japan

Green turtle nesting in Japan occurs primarily in the southern portion of the country in the Ryukyus Islands of the Okinawa prefecture and the Ogasawara Islands (AKA Bonin Islands), a group of over 30 tropical and subtropical islands (only two of which are inhabited) located approximately 1000km south of Tokyo (Kamezaki et al. 1999;

---

<sup>6</sup> Additionally, approximately 20-50 juvenile green turtles are harvested annually from waters of the Leeward Islands by poachers (Chelonia Polynesia, pers. comm.).

Horikoshi et al. 1994). The primary green turtle rookery occurs at Chichi-jima within the Ogasawara Islands which were designated a National Park by the Japanese government in 1972; most uninhabited islands have restricted access. Green turtles nest during the summer season mainly at Chichi-jima with a mean annual total of approximately 500 nesting females in recent years (Chaloupka et al. 2007).

Historically, Ogasawara was a whaling operation base and green turtles were harvested locally for their meat since the 1830s (Ishizaki 2007). Records kept since the late 19th century show a rapid decline in the sea turtle population between 1880 and 1920. By the beginning of the 20th century, efforts were made to manage marine turtles through harvest regulations and artificial hatcheries (Ishizaki 2007). Today, sea turtle harvest in the Ogasawara Islands continues under strict governmental regulation with a harvest limit of 135 mature turtles per year, alongside various resource recovery efforts led by the Ogasawara Marine Center and supported partially by the Ogasawara Village Government (Ishizaki 2007). Concurrent with regulated harvest activities, the sea turtle population has steadily increased since the early 1980's and has exhibited an estimated annual population growth rate of 6.8% per year (Chaloupka et al. 2007).

In Japan, there are eight laws and ordinances that regulate (allow via permit) or prohibit actions harmful to sea turtles, such as taking, buying, and selling turtles, their eggs, and any derivative products, or restrict access to nesting beaches. The Law for the Conservation of Endangered Species of Wild Fauna and Flora is the primary law in Japan that intends to conserve endangered species. It prohibits the capture of sea turtles and eggs for sale for all seven species and prohibits domestic assignment or transfer of endangered species listed in CITES (Umigame Hogo no tameno 2006.). This law was established in accordance with CITES and is enforced by the Japan Ministry of Environment.

## **Kiribati**

Kiribati is an island nation that consists of 32 atolls and one raised coral island that are separated into three distinct chains, the Gilbert Islands, Phoenix Islands, and Line Islands, dispersed over 3.5 million square kilometers. The westernmost islands in Kiribati (formerly known as the Gilbert or Tungaru Islands) consist of a chain of 16 atolls and coral islands including Tawara, the capital of Kiribati. In this region, green turtles have been documented nesting at Tawara Atoll, Katangateman Sandbank northeast of Makin, and another sandbank by Nonouti Island (although anecdotally, turtles have historically nested at all 16 atolls and islands except Banaba [SPC-NMFS 1979a]) with a minimum total of approximately 20 nests at Tarawa in 2007-2008 (Bell et al. 2009b). No information is available regarding nest numbers at other sites or trend for this island group.

The Phoenix Islands consist of eight low coral islands and atolls. Green turtle nesting has been observed at all eight locations including Canton, Nikumaroro, Enderbury (aka Rawaki), Phoenix, Birnie, Hull (aka Orona), Sydney (aka Manra), and McKean Islands. Canton and Enderbury Islands reportedly host the largest numbers of nesting green turtles of these eight sites. Observations in the early 1970s suggested several hundred nesting

females occurred on Canton Island (Balazs 1975b) and a survey done in the summer of 2002 recorded at least 160 old nests on Enderbury Island (Obura and Stone 2002). A combined total of 60 to 80 nests were recorded annually at the other six islands in the Phoenix group during surveys in the summers of 2000 (Stone et al. 2001) and 2002 although this is likely an underestimate of nesting activity because the peak nesting season regionally is October – November (Balazs 1995). Based on the available information, it is estimated that 100-300 green turtles may nest in the Phoenix group annually. Little to no information on trend is available for the Phoenix group.

The Line Islands consist of eleven atolls and coral islands in the central Pacific south of Hawaii, eight of which belong to Kiribati and three of which are the U.S. possessions of Palmyra, Kingman, and Jarvis. Of those under the jurisdiction of Kiribati, green turtles have been documented nesting at Christmas, Fanning, Vostok, and Caroline Islands. Information on abundance of nesting females in recent years is not available for these sites as no surveys have been conducted. Turtles appear to have declined considerably at both Fanning and Christmas Islands between the early 1800s and 1990s when human habitation began (Balazs 1995).

In Kiribati, the Wildlife Protection Act prohibits hunting, killing or capturing any wild turtle on land and fully protects the green turtle throughout the Line and Phoenix Islands (PacLII, accessed 9/10/2010). Kiribati is not a participating party to CITES.

## **Nauru**

Nauru is the world's smallest island nation and consists of one small island, approximately 21 square kilometers in area, located in the southwestern Pacific Ocean. The authors are unaware of any reports of green turtle nesting activity on Nauru. Nauru is not a participating party to CITES.

## **New Caledonia**

New Caledonia is an overseas territory of France that consists of one large Main Island (Grande Terre), the Isle of Pines, the Belep islands, the d'Entrecasteaux islands, the Chesterfields, the Loyalty Islands group, Astrolabe, Walpole, Matthew, Hunter (Fearn) islands and additional small islands and islets. The largest known nesting area for green turtles is the d'Entrecasteaux atolls, located 160 miles north of Grande Terre. It is comprised of four islands, Surprise, LeLeixour, Fabre, and Huon. There was evidence of 'major' nesting in the past, dating back to the 19<sup>th</sup> century documented by American explorer William Billings (Pritchard 1994). Pritchard (1994) described turtles to be "abundant on the southern island of Surprise, and saturation level on the additional three islands (LeLeixour, Fabre, and Huon)." Based on this survey, Pritchard (1994) estimated that 50 nesting emergences occurred per night at Huon (or approximately 2,800 nests annually). An estimate of 80 nesting females per night has also been reported (Anon. 2004). In a 1991 survey, 310 tracks were counted on Surprise island with 14 turtles tagged, 1,800 tracks and 149 turtles tagged on Huon island, 572 tracks on Fabre island, and 54 turtles were tagged in one night on both Fabre and LeLeixour islands. An additional 280 tracks and 80 nests were located on small unnamed sandy islets (Pritchard 1994). A country-wide survey of over 6,000 km of nesting habitat in December 2006 and

January 2007 identified 22 green turtle nesting locations hosting an estimated 1,000 – 2,000 nesting females annually (Limpus et al. 2009). The most recent nesting data are from a series of surveys carried out each December from 2007 to 2009 (service de la marine marchande et des pêches maritimes, 2010): on average, on all four islets, around 150 turtle tracks were counted daily in December 2007 as well as 2009. However, in 2008, more than 1,150 turtle tracks per day were reported. While trend information is not available, recent information compared to historic accounts (Pritchard 1994 & 1995a) does not suggest there has been a significant decline in abundance of green turtles nesting in New Caledonia.

Regulations related to sea turtle management in New Caledonia vary within the country. In the Loyalty Islands province (as per fishery regulations of 1985) the take of marine turtles and their eggs is prohibited from November 1 through March 31. As of January 2008, the 1985 regulations have been amended for the EEZ, the Main Island (Northern and Southern provinces), and remote islands such that it is not permitted to capture, sell, purchase, or disturb any marine turtle species or nest at anytime. Additionally, the compulsory use of handling equipment (de-hooker, line-cutter, etc.) in commercial fisheries is required for incidental catch of turtles. Regulations prohibit the export or import of marine turtles (alive or dead) or any turtle parts or products, and exceptions may be granted for customary celebrations or scientific purposes. New Caledonia is not a participating party to CITES (although France is a participating party).

## **Niue**

Niue is an island nation in free association with New Zealand. It consists of a single island approximately 256 square kilometers in area located east of Tonga. Green and hawksbill turtles occur in Niue waters (Government of Niue 2001), but the authors are unaware of any reports of green turtle nesting activity on Niue. Domestic Fishing Regulations (1996) prohibit the harvest or take of all turtle species unless approval is received from the cabinet. Niue is not a participating party to CITES.

## **Papua New Guinea**

The Independent State of Papua New Guinea (PNG) occupies the eastern half of the island of New Guinea and numerous offshore islands. Offshore islands in the northern area include New Hanover, the Tigak Islands, Djaul (including Mait Island), the St. Mathias Group (Tench, Emirau and Mussau), Tabar, Lihir, Tanga and Anir islands. Very little information is available on the abundance and trends of nesting green turtles in PNG. According to a PNG National Fisheries Authority (NFA) report, nests are raided for eggs at Nago, Atmago, and Ral islands indicating nesting activity at these locations (NFA 2007). Villagers around Kavieng indicated a peak nesting season for greens of August through October. Around Kavieng, green turtles nest at Nago Island, Atmago, Nusalaman, Usen and Lemus. In the past, green turtles used beaches on Limanak, Limalam and Nusailas Islands to nest although they are no longer in use which may be attributed to the increase in human population on these islands which led to increased harvest pressure (NFA 2007).

Long Island is a volcanic island located north of the island of New Guinea in PNG. There is limited information on green turtle nesting at this location although this rookery has been sampled for genetic analysis (Moritz et al. 2002). According to local inhabitants, greens are the most common nesters and nest all year long but with a pronounced peak nesting season from May through October (Pritchard, 1979). Nesting density was reported to be variable with anywhere from two or three nests between the villages of Malala and Point Kiau up to six or seven on a given night, according to local inhabitants. In September, 1978, Pritchard (1979) walked the seven mile stretch of beach between the villages of Malala and Point Kiau on Long Island and observed twelve nesters and tracks of at least seven others in one night. He estimated a total of 35 for the night with more likely beyond the scope of the survey (Pritchard 1979). Informants revealed that 20 to 30 nesting turtles per month were eaten by island residents and also that, at the time, turtles were considered “as plentiful as they ever have been.” Much of this information is inconsistent making it difficult to estimate an abundance of annual nesting females at this site. Harvest pressure was apparent in 1978 and has likely continued so a more recent survey of turtle nesting activity on Long Island is needed for a reliable current estimate of nesting females.

In January 2003, the first assessment of turtle stocks in the Milne Bay Province commenced at Panayayapona Island of the Brooker Islands (Kinch 2003a). Sixteen green turtle nests were documented from January 21-27, 2003 with a total of 71 tracks recorded. For comparison, during a reconnaissance survey on Nov 28, 1998 a total of 177 tracks (not discerned by species) were recorded at Panayayapona. The previous year 126 tracks were counted (not discerned by species). In mid-December 1997 an average of 30 to 40 turtles arrived each night to nest and on one night in mid-January 2002, 72 tracks (not discerned by species) were counted (Kinch 2003b). More intensive surveying December 1-21, 2004 at the Jomard Islands (Panadaludalu and Panayayapona), Siva of the Bramble Haven group, and Irai, Pananiu, and Tobiki islands of the Conflicts group resulted in 115 green turtle nests recorded (Wangun et al. 2004).

A comprehensive survey of PNG for green turtle nesting abundance has not been done nor is current trend information available, but previous (dated) studies indicated that numbers of green, hawksbill, and leatherback turtles were decreasing throughout PNG (Pritchard 1982; Spring 1982; Bedding and Lockhart 1989). It is likely that this declining trend has continued over time, with the exception of green turtles nesting in areas of Seventh Day Adventists (this religion prohibits eating meat, including turtles), such as Mussau and Emirau Islands in the St. Mathias Group (NFA 2007). In these areas, Pritchard (1995) reports a “noticeable increase in the turtle populations over a 30 to 50 year period,” although NFA (2007) reports that people from Kavieng and Manus visit the islands to harvest turtles.

In PNG, marine resources and lands are owned by a large number of clan and sub-clan groups whose tenure rights are recognized in the national Constitution. With respect to sea turtles, the 1976 Fauna (Protection and Control) Act restricts the harvesting of protected wildlife, the devices and methods by which fauna may be taken, and the establishment of localized protective regimes on land and waters under customary tenure

(Kinch 2006). Additionally the 1979 International Trade Act regulates and restricts the export of CITES listed species. In PNG, only leatherback turtles are protected under the Fauna (Protection and Control) Act that makes killing of leatherbacks or taking of leatherback turtle eggs illegal with fines of 500-1000 kina (100 to 300 USD). Any person who buys or sells or offers for sale, or has in possession leatherback turtle eggs or meat can also be fined 500 kina. The Act does not formally protect green turtles and makes provisions for persons with customary rights to take or kill turtles, but states that turtles cannot be taken, killed, or sold during the months of May through July. Furthermore, the Act stipulates payments for turtles: (a) K20.00 for a turtle less than 60 cm in length; and (b) K30.00 for a turtle of 60 cm or more in length. The PNG government Department of Environment and Conservation has the authority and responsibility to enforce laws and environmental Acts. PNG is a participating party to CITES.

### **Pitcairn Islands**

The Pitcairn Islands are an overseas territory of the United Kingdom (UK) consisting of four volcanic islands. Green turtles nest at Henderson Island with an estimated total of 10 females annually (Brooke 1995). No nesting was recorded at Pitcairn, Ducie, or Oeno Islands during the 1991-1992 nesting season. Pitcairn and Ducie were deemed to have unsuitable substrate for nesting while Oeno had suitable substrate but no activity was observed (Brooke 1995). This small nesting assemblage does not appear to be threatened by direct harvest or other major anthropogenic sources of impact. As per local Government Regulations (1982), no person may harass, hunt, kill or capture any sea turtle (*Cheloniidae* and *D. coriacea*), and exception may be granted under permit for scientific purpose or for traditional subsistence use (SPREP 2007). As a territory of the UK, the Pitcairn Islands are subject to the wildlife conservation laws of the UK which is a participating party to CITES.

### **Samoa**

The Independent State of Samoa consists of two large islands, Upolu and Savai'i, and eight small islets and is located west of American Samoa in the south Pacific. Although both green and hawksbill turtles are known to occur in the waters of Samoa, only hawksbill turtles nest here with no green turtle nesting reported (Witzell 1982). Adult greens have been observed near reefs year-round, and from December through February have been observed gathering near reef passages connecting large lagoonal foraging areas near Upolu Island. Witzell (1982) surmised that these adults may be part of the group that nests on Rose Atoll during August-September.

Local Fisheries Regulations in Samoa prohibit fishing for, possession, or sale of greens and hawksbills under 70cm (27.6 inches) curved carapace length as well as the disturbance or take of nests or eggs. The Ministry of Agriculture and Fisheries (formerly the Department of Agriculture, Forests, and Fisheries) is the responsible authority to manage fishery resources, including sea turtles, and enforce local fisheries regulations in Samoa. Additionally, the Marine Wildlife Protection Regulations (2009), under the Ministry of Natural Resources and Environment, make exemptions for subsistence take of turtles, prohibit captivity of turtles (unless permitted), prohibit the commercial capture and sale of turtles, protect turtles and eggs during the nesting season (November to



February), prohibit the sale, purchase and possession of eggs, and require that any turtle caught during fishing activities be released and reported. Samoa is a participating party to CITES.

## **Solomon Islands**

The Solomon Islands are located east of Papua New Guinea and consist of nearly one thousand islands. More is known about hawksbill nesting in the Solomon Islands with limited information available regarding current overall nesting of green turtles. McKeown (1977) estimated that 45 green turtle nests were laid in the Arnavon Islands that year. The Solomon Islands Fisheries Division undertook the first extensive turtle survey in the Solomons focused primarily on the Arnavon Islands in Isabel Province (Vaughan 1980 and 1981). Vaughan (1981) estimated that the number of breeding individuals of all sea turtle species combined in the Solomons was about 1,500 and that 42% of hawksbill and green turtles present in the Solomon Islands nested in Isabel Province. This survey recorded 53 green turtle nests on Kerehikapa, Arnavon Island, during the months of September to March, with Hakelake Island also supporting 15 to 20 nests per year. Ramohia and Pita (1996) identified only five green turtles nesting in the Arnavon Islands during summer of 1995. Vaughan (1981) also documented green turtle nesting activity within the provinces of Choiseul, Shortlands and Makira primarily on the islands of Wagina, Ausilala, Maifu, Balaka, and Three Sisters (Malaulaul and Malaupaina), with approximately 50 to 100 green turtle nests laid per year at each island. While Vaughan's 1980 survey noted anecdotal reports of a reduction in abundance due to high harvest pressure<sup>7</sup> on nesting females and their eggs, Leary and Laumani (1989) estimated a modest increase of nesting activity in Isabel province of 259 to 438 nests, compared to 211 to 341 nests in 1980 (not including the Arnavon Islands). This discrepancy is likely a result of normal fluctuations in turtle nesting activity and not indicative of a measurable trend in this nesting assemblage. Long distance migrations between foraging and breeding grounds have been documented indicating that there is reciprocal movement between Australia and the Solomon Islands (Broderick, unpublished 1998).

Currently, the only continuous nesting beach monitoring projects in the Solomons occur at the Arnavons (green and hawksbill), Sasakolo (leatherback), and Tetepare (leatherback and green) (C.Siota pers. comm.). Updated information from the Arnavons and Tetepare for green turtles was not available at the time of printing. Sufficient data on abundance and trend for Solomon Islands green turtles are not available.

The Solomon Islands Fisheries Act (1993) regulations prohibit the sale, purchase, or export of sea turtle species or their parts, protect nesting turtles and eggs during the breeding season (June to August and November to January), and contain specific protections for leatherback turtles (SPREP 2007). The Solomon Islands is a participating

---

<sup>7</sup> Based on intensive surveying between 1993 and 1996, Broderick (unpublished, 1998) concluded that the Solomons serve as important developmental habitat for juvenile green turtles, but that a large proportion of turtles are being harvested. Pita and Broderick (2005) estimated that over 1,000 green turtles are harvested per year in the Solomons in three different villages (Kia, Wagina, Katupika) of Isabelle and Choiseul provinces.

party to CITES and the Wildlife Protection and Management Act (1998) prohibits the export of five turtle species or their derivative products (greens, hawksbills, loggerheads, olive ridleys, and leatherbacks).

## **Tokelau**

Tokelau consists of three coral atolls, Atafu, Nukunonu, and Fakaofu, all of which are known to have green turtle nesting. Balazs (1983b) estimated 120 total nesting females annually in Tokelau. Sea turtle capture rates declined from the early 1900s to the 1980s despite more sophisticated hunting methods, indicating a likely population decline (Balazs 1983b). Updated information regarding abundance and trends of nesting green turtles in Tokelau was not available to the authors at the time of printing.

According to a 1998 marine resources survey at Fakaofu, the local council of elders has established village rules stating that when a turtle is caught, it must be shared among the village using a traditional system or resource sharing called Inati (Passfield 1998). Local village rules also protect sea turtles while they are nesting. Tokelau is not a participating party to CITES.

## **Tonga**

The Kingdom of Tonga is composed of at least 170 islands, 36 of which are inhabited. Islands are grouped into three main regions: the Ha`apai Group, Vava`u Group, and Tongatapu Group. Nesting takes place from October to February with peak nesting in December and January (Bell et al. 1994). Green turtles nest in low levels on several islands in the Ha`apai Group as well as islands in the Vava`u Group, with an estimated 10 to 20 green turtle nests annually based on anecdotal information from turtle hunters (Havea and MacKay 2009). Sporadic nesting surveys were carried out in the Ha`apai Group in December 1971, December 1973, and December 2007 to January 2008 (Bell et al. 2009c) although most did not distinguish between hawksbill and green nests and effort was not consistent among surveys. Wilkinson (1979 in Bell et al. 1994) reported that fishermen have seen green turtles nesting on the uninhabited islands of Nukufaiva, Fetoa, Mango, and Malinoa in the Ha'apai group but nesting numbers were not provided. Based on limited available information from past surveys, it appears that 10 to 100 females may nest annually in Tonga, with numbers likely toward the lower end of that range.

Havea and MacKay (2009) surveyed fishermen for their perceptions of sea turtle abundance trends in the Ha`apai group. In spite of previous reports and an apparent decline in nesting turtles, less than 50 percent of fishermen reported that turtle stocks were declining and almost 40 percent indicated stocks were increasing. However, the survey did not distinguish between greens and hawksbills. Directed take of green turtles for consumption and sale still occurs in Tonga and laws are generally not adhered to or enforced (Havea and MacKay 2009). Limited available data on nesting in addition to these survey results suggest there may be a decline in green turtle nests (Havea and MacKay 2009).

Fisheries Conservation and Management Regulations (1994) prohibit the possession, disturbance, take, sale, purchase, or export of turtle eggs, sale, purchase, or export of

hawksbills or their shells, and use of a spear gun to take a turtle and establish closed seasons for leatherback turtles January 1 to December 31 (Bell et al. 1994; Folumoetui'i, 2006). Hawksbills are the only sea turtle species fully protected via Tongan legislation and harvest of other turtle species is permitted seasonally (November to February) with a minimum size specified (shell length of <45cm) (Folaumoetu'i 2006). Tonga is not a participating party to CITES.

## **Tuvalu**

Tuvalu is an independent nation made up of nine coral islands and atolls. Green turtles nest in the capital of Funafuti as well as on several outer islands (Pita 1979). The only available information on nesting turtles in Tuvalu is from a 10 day survey of nesting sites on Funafuti conducted in December 2006 where a total of nine nest sites were identified (Alefaio et al. 2006). In 1979, turtle meat was rarely consumed in the capital of Funafuti but turtles were still taken from the water and nesting beaches for consumption in the outer islands where there was no refrigeration (Pita 1979). According to interviews with local fishermen in 2006, the number of turtles sighted and harvested has declined rapidly (Alefaio et al. 2006).

In Tuvalu, the Wildlife Conservation Ordinance (1975) prohibits hunting, killing or capturing any wild turtle on land except under and in accordance with the terms of a valid written license granted to that person by the Minister (Government of Tuvalu 1975). Tuvalu is not a participating party to CITES.

## **Vanuatu**

Vanuatu consists of approximately 82 islands, 65 of which are inhabited. Turtles in Vanuatu are described as "plentiful" with Malekula island identified in 1979 as an important nesting area with 40 to 120 turtles nesting annually (although species was not specified, this likely refers to a combination of greens and hawksbills) (Pritchard 1982 in Pritchard 1995a). Currently, the only published information on sea turtle nesting activity is summarized in Petro et al. (2007) based on interviews of knowledgeable turtle monitors and limited surveys that occurred from November to December 2002 and January to February 2003, focused primarily on leatherback turtles. During a survey at Votlo, Southern Epi Island, two green turtles were tagged and 10 false crawls and 15 nests were recorded. Current information collected at Wan Smolbag workshops in 2007 and 2008 by monitors of the Vanua-Tai network identified over 189 nesting sites on 33 islands of Vanuatu, with approximately 200 turtles (both green and hawksbill) nesting at Malekula island per year (Fletcher and Petro, unpublished 2009). Additionally, Santo Island and its offshore neighboring island of Thion support 50 or more nesting turtles per year, and approximately 30 turtles nest annually at Tegua and Hiu islands. Coverage of Vanuatu's beaches is not yet comprehensive so total nesting activity may be underestimated. A number of sites have emerged over the past few years as potential index sites, in particular the Bamboo Bay area on the island of Malekula, and possibly Moso and Wiawi (G. Petro pers. comm.). Current trend information is not available for green turtles in Vanuatu. Green turtles and their eggs are commonly harvested and there was recent momentum to revive traditional management systems to regulate (or sustainably manage) community-based harvest of turtles (Hickey 2007). Primary threats

to green turtles identified in Vanuatu in addition to direct harvest include nest predation by dogs, wave inundation and beach erosion.

Fisheries Regulations under the new Vanuatu Fisheries Act (2009) prohibit the take, harm, capture, disturbance, possession, sale, purchase of or interference with any turtle nest (or any turtle in the process of nesting), and the import, or export of green, hawksbill, and leatherback turtles or their products (shell, eggs, or hatchlings). The Act also prohibits the possession of turtles in captivity. A person may apply in writing to the Director of Fisheries for an exemption from all or any of these provisions for the purposes of carrying out customary practices, education, and/or research. Vanuatu is a participating party to CITES.

### **Wallis and Futuna**

Wallis and Futuna consist of three main islands and low coral or volcanic islets. The authors are unaware of any reports of green turtle nesting activity on any of the islands or islets at Wallis and Futuna. Wallis and Futuna is not a participating party to CITES.

## Discussion and Recommendations

This is the first comprehensive compilation of peer-reviewed publications, gray literature, and local government reports of green turtle nesting information from locations throughout Oceania. A total of 189 sites were identified (Table 1 and Figure 1), but many more nesting sites are likely to occur throughout the region as we describe only those that have been surveyed, monitored, or observed and reported in publicly available documents covering the past 30 to 40 years. A historical perspective or baseline of nesting locations and abundance in the region (e.g., 200 to 300+ years ago)<sup>8</sup> is beyond the scope of this document but is necessary to identify recovery goals for green turtles in Oceania.

Although there are currently a number of active monitoring projects throughout the Pacific (Figure 2), consistent long-term datasets necessary to draw reliable conclusions about trends of annual nester abundance only exist for four of the 189 sites in the region: Hawaii (Chaloupka et al 2008), Japan, and two sites in Australia (Chaloupka 2007). According to our review, a number of sites in addition to these four are thought to have relatively substantial nesting activity (between 100 and 1,000 nesting females annually; Table 1). The varied levels of survey effort and methodology within and among countries from year to year, and discrepancies in types of data reported for many sites make it difficult to clearly discern nesting female abundance and trends. With respect to management objectives, there is an emergent and immediate need to initiate or facilitate consistent long-term monitoring projects at more locations throughout the region. The potential for such projects to be established (or maintained), considering cultural, financial, and logistical limitations, should be investigated as additional index sites would allow scientists and managers to more adequately assess overall green turtle abundance, stock structure, trends, and threats in the Pacific. Areas for consideration include: Helen Reef in Palau, Ulithi Atoll<sup>9</sup> in Yap, d'Entrecasteaux Reef complex in New Caledonia, Scilly Atoll<sup>10</sup> in French Polynesia, Malekula Island<sup>11</sup> in Vanuatu, and the Coringa-Herald National Nature Reserve in the Coral Sea. Additional sites reported to support, or to have once supported, significant nesting activity that do not have consistent survey efforts in place include: Rose Atoll of American Samoa, Bikar and Erikub Atolls in the Marshall Islands, Canton Island in the Phoenix Islands of Kiribati, and Palmerston Atoll in the Cook Islands. Many of these areas are extremely remote and logistically challenging to access and/or may involve associated cultural challenges and island-specific politics that may affect beach access and approvals necessary to conduct monitoring activities. Rapid

---

<sup>8</sup> Baseline historical information from naval, explorer, economic, ethnographic, and anthropological sources dating from the 1600s to present is currently in preparation by the NMFS PIFSC Marine Turtle Assessment Program.

<sup>9</sup> NMFS PIRO has supported monitoring activities at Ulithi Atoll with funding and technical expertise since 2005 and, as available funding permits, intends to continue support for at least 10 years.

<sup>10</sup> Scilly Atoll has not been surveyed since 1995, however, a local NGO, Chelonia Polynesia, with funding support from NMFS PIRO plans to revisit this atoll in 2010-11 to gather updated census information on this nesting population and collect genetic samples.

<sup>11</sup> Green turtle monitoring activities in Vanuatu are led by the WanSmolbag Theater Turtle Conservation Program. WanSmolbag has received support from NMFS Southwest Region since 2007 for monitoring activities of the Vanua-tai turtle network, and in 2010 received additional support from NMFS PIRO to incorporate genetic sample collection in their monitoring protocol.

assessments of nesting activity occurring during peak nesting periods (if known) over consecutive years (each conducted within a 3 to 4 week timeframe) may be an option to assess nesting activity to inform management at sites that can be accessed (i.e., permissions granted).

A cursory look at the relationship between local legislation and nesting abundance indicates a slight pattern (Figure 1), and although this should be interpreted with caution, we confirm previously described observations (Bjorndal 1982 revised in 1995; Eckert 1993). In general, larger nesting assemblages occur in countries that have full or relatively strong legal protection for sea turtles while countries with weaker or less effective protective regulations in place tend to support smaller nesting populations or those that are not consistently monitored and assumed to be small. Additionally, larger nesting assemblages appear to occur in countries that are more developed and have advanced science and conservation programs that can produce information on marine turtle resources on a regular and consistent basis. In contrast, smaller, less developed countries likely have fewer resources for scientific research and/or alternative priorities. Sea turtles are also highly migratory and those that nest in one region are vulnerable to threats or causes of mortality in other geographic areas, often with different levels of legislative protection, during different phases of development and migration (Craig et al. 2004; Dutton and Squires 2008; Seminoff 2004). For example, Australia has relatively strong legislative protections (Figure 1), yet tag return data confirm that Australian turtles are captured in the Torres Strait, New Guinea, and Solomon Island waters (Limpus et al. 2003; Limpus 2009) where the same level of protective legislation does not exist. Similarly, post-nesting females from French Polynesia, the Cook Islands and Rose Atoll have been recorded traveling westward, primarily to Fiji (Craig et al. 2004), passing through several EEZs and high seas areas where various domestic and international fisheries operate, highlighting the value of collaborative management strategies among nations.

Where indigenous and unregulated subsistence harvest of sea turtles is permitted, many of the existing regulations in place are not appropriate because they do not reflect a basis in the biology of sea turtle life history. Rather, they appear to mimic regulations designed to optimize population levels of fish which often have very different life history characteristics than turtles. Minimum size regulations designed to protect small juveniles yet permit the harvest of large juveniles (subadults) and adults do not provide the protection necessary for long-term sustainable management of sea turtle populations as large juveniles and adults have the highest reproductive value of all life stages (Crouse, 1999; Heppell 1998). Throughout Micronesia, Melanesia, and some parts of Polynesia (Figures 1, 1a, and 1b), there are many nesting sites without reliable estimates of annual nesting females or with low numbers of nesting females (less than 100) where turtles are subject to biologically irrelevant regulations (i.e., allowable size limits, harvest open/closed seasons). Micronesia in particular is reported to support a large number of known rookeries yet hosts few monitoring projects and imposes regulations that are inappropriate for effectively managing sea turtle populations. As such, Micronesia is still (as previously suggested by Pritchard 1977, Buden and Edward 2001, and NMFS and USFWS 1998) considered a priority area to initiate more monitoring activities coupled

with intensive and integrated education and awareness efforts. Government collaboration to develop stronger, more biologically relevant regulations for the effective management of local sea turtle populations is warranted.

Many countries throughout Oceania are signatories or participating parties in various regional and international arrangements (Appendix I). Nine of the 22 countries in the region are participating parties to the Convention on International Trade in Endangered Species (CITES) which prohibits all commercial trade of marine turtles and marine turtle derived products. Several countries throughout the region are also participants in other global environmental agreements including: the Convention on Biological Diversity (CBD), Convention on Migratory Species (CMS), and the Ramsar Convention on Wetlands. In total, there are eight regional arrangements in Oceania that address aspects of marine turtle conservation, protection, or management. The three most prominent arrangements include the Western and Central Pacific Fisheries Commission (WCPFC), Pacific Islands Forum Fisheries Agency (FFA), and Secretariat of the Pacific Regional Environment Programme (SPREP). All countries discussed in this technical memorandum are members of SPREP (with the exception of Japan) and SPREP has developed a non-binding 2008-2012 Marine Turtle Action Plan (MTAP) (SPREP 2008). The MTAP identifies actions focusing on reducing threats including unsustainable harvest, nest predation, incidental capture in commercial fishing, degradation of habitat, pollution, marine debris, pathogens, boat strikes, and climate change. It also highlights national and regional priority actions that have been identified and endorsed by member countries for marine turtle conservation and monitoring work in the region. Therefore, a foundation exists for international collaboration in the Pacific to address the management and conservation challenges discussed above pertaining to marine turtles and regulations governing subsistence or direct harvest.

The vast majority of the 189 sites described here have 100 or fewer documented annual nesting females, with many likely to have 10 or less (Table 1). While these nesting assemblages make up a small proportion of the overall Oceania green turtle nesting aggregation, they may possess unique adaptations in genetic diversity and ecological significance to their particular environments, and may therefore be important for recovery of the species (Bjorndal and Bolten 2008). There is legitimate concern over the loss of small isolated nesting aggregations and the implications for the species as a whole, as well as associated consequences for local ecosystems (McClenachan et al 2006; Bjorndal and Jackson 2003). Female green turtles tend to exhibit strong natal homing (Miller 1997), returning to the beach from which they hatched to lay their eggs, making particular rookeries reproductively isolated over ecological timescales. Although 'mistakes' in natal homing occur with sufficient frequency to facilitate founder or seed populations and some genetic exchange between conspecific rookeries over short evolutionary timescales, the decline or loss of a given rookery is not likely to be compensated for by natural recruitment of females hatched elsewhere (at least over ecological timescales relevant to immediate human conservation interests) (Awise and Bowen 1994).

We recognize that this synthesis of information refers to nesting females which is just one aspect of sea turtle life history affecting population dynamics. An understanding of key foraging habitats, foraging population demographics (including males, juveniles and subadults), and mixed stock genetic analyses at foraging grounds are necessary for a complete assessment of population status and sources of mortality (as described and summarized in NMFS and FWS 2007, and NRC 2010). While there are indications that some Pacific Island nations may provide important green turtle foraging habitats (Balazs and Forsyth 1986; Balazs and Chaloupka 2005; Chaloupka and Limpus 2001; Craig et al. 2004; Kolinski 2004; Meylan 1982; Pritchard 1995a,b; Pultz et al. 1999), a region-wide understanding of sea turtle foraging, stock connectivity, and the relative contribution of Pacific stocks within foraging habitats does not currently exist. The genetics lab at the NOAA Fisheries Service Southwest Fisheries Science Center continues to work with partners like the University of Canberra in Australia to sufficiently characterize Pacific green turtle nesting stocks with informative genetic markers in order to improve the ability to assign stock origin of individual animals in foraging and migratory habitats. In the future, this genetic information may be applied to determine Regional Management Units (such as those proposed by Wallace et al. [*in press*]) or to evaluate the need for Distinct Population Segments as recognized by the ESA.

In summary, more information is available for green turtle nesting in Oceania than has been previously summarized or compiled, but the available information suggests there are large gaps in current information gathering and monitoring efforts. This compilation is a step in identifying those gaps and will assist scientists and managers in prioritizing projects and funding considerations in accordance with where needs may exist to bolster management and assessment efforts for recovery planning. In addition to research and monitoring, local legislation in countries with green turtle nesting assemblages should be evaluated and government collaborations encouraged to facilitate implementation of more biologically effective regulations coupled with integrated education and awareness initiatives. These efforts will help prioritize and direct future research, management activities, and international collaborations to advance green turtle conservation and recovery efforts in the Pacific.



## **Acknowledgments**

This work would not have been possible without the involvement and generosity of our friends and colleagues at SPREP: Jeff Kinch, Lui Bell, Anne Trevor, and Paul Anderson. Numerous additional in-country associates provided reports, grey literature, and information, and assisted with document review: George Petro, Michelle Fletcher, Catherine Siota, Sebastien Goutenegre, Yoshimasa Matsuzawa, Kirstin Dobbs, Shawn Wusstig, Alden Tagarino, Patrick Opay, Stacy Hargrove, Tammy Summers, John Pita, Joshua Eberdong, and Jennifer Cruce. George Balazs and the PIFSC Marine Turtle Research Program also provided numerous documents from their archives as well as document review. We thank Kevin Kelly for his comments and editorial assistance. This document was in part inspired by the WIDECAST 2007 Atlas of Sea Turtle Nesting Habitat for the Wider Caribbean Region (Dow et al. 2007) and we therefore acknowledge Wendy Dow, Karen Eckert and other co-authors.



## Tables and Figures

**Table 1:** Summary of Information for Nesting Green Turtles in Oceania Including Current Generalized Legislation

Country/Territory/State	Est. # Known Sites	Est. # Annual Nesting Females (min.)	Est. Trend	Legislation
American Samoa (USA)	3	30 - 100	Unknown	A
Australia (NGBR)	5	10,000-25,000	Stable*	B
Australia (SGBR & HNNR)	21	5,000 – 10,000	Increasing* (SGBR)	B
Cook Islands	11	<10	Unknown	D
CNMI (USA)	16	<10	Unknown	A
Federated States of Micronesia	16	530 - 1,300	Unknown	C
Fiji	2	10 - 100	Decreasing	A
French Polynesia	7	300 - 400	Unknown	B
Guam (USA)	15	<10	Unknown	A
Hawaii (USA)	12	400 - 500	Increasing*	A
Japan (Ogasawara Isl.)	1	400 - 500	Increasing*	A
Kiribati	15	80 - 800	Unknown	C
Nauru	0	0	NA	D
New Caledonia	4	1,000 - 2,000	Stable	A <sup>^</sup>
Niue	0	0	NA	A
Papua New Guinea	15	10 - 100	Unknown	D
Pitcairn Islands	1	<10	Unknown	E
PRIAs (USA)	2	<10	Unknown	A
Republic of Marshal Islands	9	300 - 400	Decreasing	C
Republic of Palau	13	250 - 400	Unknown	C
Samoa	0	0	NA	C
Solomon Islands	8	80 - 700	Unknown	C
Tokelau	3	<10	Unknown	D
Tonga	3	<10	Unknown	C
Tuvalu	1	<10	Unknown	C
Vanuatu	6	20 - 100	Unknown	A
Wallis and Futuna	0	0	NA	E
<b>Total</b>	<b>189</b>	<b>18,480 – 42,470</b>		


\* Chaloupka et al. (2007)

\*\* Generalized Legislation Categories (excluding reference to CITES, international arrangements (Appendix 1) and commercialized export of sea turtles):

- A. Allowable (permitted/regulated) take (includes moratoriums)
- B. Allowable (permitted/regulated) take, and recognized (unregulated) indigenous/subsistence harvest (i.e., protections afforded via indigenous cultural practices)
- C. Open/closed harvest periods, or minimum size limits, or protected when on land only (i.e., nesting females & nests), and/or recognized indigenous/subsistence harvest
- D. No legislation specific to green turtles (but may include loose protections of nesting turtles when on land), and recognized indigenous/subsistence harvest
- E. Sea turtle legislation unclear or unknown to authors

<sup>^</sup> Most of New Caledonia falls into category A however the Loyalty Islands have an open harvest season for sea turtles (category C).







**Figure 1**

**Green Turtle Nesting and Legislation in Oceania**

This map shows nesting sites for green turtles throughout Oceania and the estimated minimum annual nesting females at each site along with the varied categories of legislation associated with each country where nesting takes place.





Geographic Coordinate System: WGS84



0 500 1,000 2,000 3,000 4,000 Kilometers

Map by: Kim Matson, NOAA Fisheries Pacific Islands Regional Office, August 2010







Figure 2

**Green Turtle Nesting Beach Projects**

This map shows the locations of active monitoring projects at green turtle nesting beaches throughout Oceania that have collected data within the past five years (since 2005).

Geographic Coordinate System: WGS84



Map by: Kim Maison, NOAA Fisheries Pacific Islands Regional Office, August 2010







## Appendix I: Pacific Countries and Existing International Arrangements

Pacific Countries and Existing International Arrangements for Sea Turtle Conservation, Protection and Management (information compiled for and presented at 20<sup>th</sup> SPREP meeting, November 19-20, 2009, Apia, Samoa)

Country	CBD	CITES	CMS	CTI	FFA	IAC	IATTC	IOSEA	IOTC	RAMSAR	SPREP	WCPFC
<i>American Samoa</i>											S	P
<i>Australia</i>	P	P	P		S			S	P	P	S	P
<i>Cook Islands</i>	P		P		S		C				S	P
<i>FSM</i>	P				S						S	P
<i>Fiji</i>	P	P			S					P	S	P
<i>French Polynesia</i>											S	P
<i>Japan</i>	P	P					P		P	P		P
<i>Kiribati</i>	P				S						S	P
<i>Marshall Islands</i>	P				S					P	S	P
<i>Nauru</i>	P				S						S	P
<i>New Caledonia</i>											S	P
<i>Niue</i>	P				S						S	P
<i>Palau</i>	P	P	P		S					P	S	P
<i>PNG</i>	P	P		S	S					P	S	P
<i>Samoa</i>	P	P	P		S					P	S	P
<i>Solomon Islands</i>	P	P		S	S						S	P
<i>Tokelau</i>					S						S	P
<i>Tonga</i>	P				S						S	P
<i>Tuvalu</i>	P				S						S	P
<i>USA</i>	S	P				P	P	S		P	S	P
<i>Vanuatu</i>	P	P			S		P		P		S	P
<i>Wallis and Futuna</i>											S	P

Party or Participating Entity/Territory = P

Signatory = S

Cooperating non-Member = C

### Appendix 1 Acronyms:

**CBD**: Convention on Biological Diversity; **CITES**: Convention on International Trade in Endangered Species; **CMS**: Convention on Migratory Species; **CTI**: Coral Triangle Initiative on Coral Reefs, Fisheries, and Food Security; **FFA**: Forum Fisheries Authority; **IAC**: Inter-American Convention for the Protection and Conservation of Sea Turtles; **IATTC**: Inter-American Tropical Tuna Commission; **IOSEA**: Indian Ocean South-East Asian Sea Turtle Memorandum of Understanding; **IOTC**: Indian Ocean Tuna Commission; **Ramsar**: Ramsar Convention on Wetlands of International Importance; **SPREP**: Secretariat of the Pacific Regional Environment Program; **WCPFC**: Western/Central Pacific Fisheries Commission.



## References

- Allen, M.S. 2007. Three millennia of human and sea turtle interactions in remote Oceania. *Coral Reefs* 26: 959-970
- Alefaio, S., T. Alefaio, and A. Resture. 2006. Turtle monitoring on Funafuti, Tuvalu December 4<sup>th</sup>-14<sup>th</sup> 2006. Report of Survey administered by the Institute of Marine Resources, the University of the South Pacific, Suva, Fiji. 8pp.
- Anonymous. 2004. Announcement of Green Turtle Research Project, d'Entrecasteaux Reef, New Caledonia. *Marine Turtle Newsletter* 103:25.
- Awise, J. C. and B.W. Bowen. 1994. Investigating sea turtle migration using DNA markers. *Current Opinions in Genetics and Development*; v. 4 (6): 882-886.
- Balazs, G.H. 1975a. Green turtle uncertain future: Protection vial if remnant population is to survive. *Defenders*; v. 50(6): 521-523.
- Balazs, G.H. 1975b. Marine turtles in the Phoenix Islands. Atoll Research Bulletin No. 184.
- Balazs, G.H. 1976. Green turtle migrations in the Hawaiian archipelago. *Biological Conservation* 9:125-140.
- Balazs, G. H. 1980. Synopsis of biological data on the green turtle in the Hawaiian Islands. NOAA Tech. Memo. NOAA-TM-NMFS-SWFC-7.
- Balazs, G.H. 1983a. Subsistence use of sea turtles at Pacific Islands under the Jurisdiction of the United States. Southwest Fisheries Center, Honolulu Laboratory, Administrative Report H-83-17.
- Balazs, G.H. 1983b. Sea turtles and their traditional usage in Tokelau. *Atoll Research Bulletin* 279:1-30
- Balazs, G.H. and R.G. Forsyth. 1986. Status and ecology of marine turtles at Johnston Atoll: 1985 Assessment. NMFS SWFC Honolulu Laboratory Administrative Report H-86-9.
- Balazs, G.H. 1995. Status of sea turtles in the central Pacific Ocean. In: Bjorndal, K. (ed.). *Biology and Conservation of Sea Turtles*. Smithsonian Institution Press. Washington D.C., 615 pp.
- Balazs, G.H., P. Siu and J.P. Landret. 1995. Ecological aspects of green turtles nesting at Scilly Atoll in French Polynesia. In: Richardson, J.I. and T.H. Richardson (compilers). *Proceedings of the Twelfth Annual Workshop on Sea Turtle Biology and Conservation*, NOAA Technical Memorandum NMFS-SEFSC-361, p.7-10.
- Balazs, G.H. and M. Chaloupka. 2004. Thirty-year recovery trend in the once depleted Hawaiian green sea turtle stock. *Biological Conservation*, 117:491-498.
- Balazs, G.H. and M. Chaloupka. 2005. Spatial and temporal variability in somatic growth of green sea turtles (*Chelonia mydas*) resident in the Hawaiian Archipelago. *Marine Biology* 145: 1043-1059.
- Balazs, G.H. 2009. Historical summary of sea turtle observations at Rose Atoll, American Samoa, 1839-1993. Unpublished internal report compiled by George Balazs, Marine Turtle Research Program, NOAA, National Marine Fisheries Service, Pacific Islands Fisheries Science Center.
- Barr, J. M. 2006. Community based sea turtle monitoring and management at Helen Reef, Hatohebei State, Republic of Palau. Master's Thesis, Oregon State University. 216p.

- Batibasaga, A. 2002. Sea turtle status & conservation initiatives in Fiji. In: I.Kinan (ed.) Proceedings of the Western Pacific Sea Turtle Cooperative Research & Management Workshop. WPRFMC. Feb 2-5, 2002.
- Batibasaga, A., S. Waqainabete, & A. Qauqau. 2006. Notes on Fijian sea turtles: estimates on population status. Fiji Fisheries Department, PO Box 3165, Lami, Fiji. Information provided for Sea Turtle Working Group Meeting – Nadave / CATD 31st May – 1st June 2006.
- Bedding, S. and B. Lockhart. 1989. Sea turtle conservation emerging in Papua New Guinea. *Marine Turtle Newsletter* 47:13.
- Bell, L., J. Ward, and P. Ifopo. 2009a. Marine turtles strand on land after the Samoa Tsunami. SPREP/DEC technical report.
- Bell, L.A.J., N. Ruata, and R. Bebe. 2009b. Kiribati Marine Turtle Profile. Unpublished draft report for the Secretariat of the Pacific Regional Environment Programme.
- Bell, L.A.J., L. Matoto, and `U. Fa`anunu. 2009c. Project Report: Marine Turtle Monitoring Programme in Tonga. Marine Turtle Conservation Act Project Report. 15 pp. plus appendices.
- Bell, L.A.J., U, Fan`anunu, and T. Koloa. 1994. Fisheries resources profiles: Kingdom of Tonga. FFA report 94/05. Available from: <http://www.sprep.org/att/IRC/eCOPIES/Countries/Tonga/7.pdf>.
- Bjorndal, K.A. 1982 (revised 1995). Biology and Conservation of Sea Turtles, revised edition. Smithsonian Institution Press. Washington D.C., 615 pp.
- Bjorndal, K. A. and J.B.C. Jackson. 2003. Roles of sea turtles in marine ecosystems: reconstructing the past. In: Lutz, P. L. ,Musick, J. A. ,Wyneken, J. (editors), The Biology of Sea Turtles. Volume II. CRC Marine Biology Series, CRC Press, Inc.: Boca Raton, London, New York, Washington D.C. 455 pp.; p. 259-273.
- Bjorndal, K.A. and A. B. Bolten. 2008. Annual variation in source contributions to a mixed stock: implications for quantifying connectivity. *Molecular Ecology*; 17; 2185–2193.
- Broderick, D. unpublished, 1998. Subsistence Harvesting of Marine Turtles in the Solomon Islands. Patterns of resource use in Kia, Wagina and Katulika communities, Isabel and Choiseul Provinces.
- Brooke, M. de L. 1995. Seasonality and numbers of green turtles *Chelonia mydas* nesting on the Pitcairn Islands. *Biological Journal of the Linnean Society* 56:325-327.
- Buden, D.W. and A. Edward. 2001. Abundance and utilization of sea turtles on Pohnpei, Federated States of Micronesia: Islander’s Perceptions. *Micronesica* 34(1):47-54
- Campbell, L.M. 2003 Contemporary Culture, Use, and Conservation of Sea Turtles. In: P.A. Lutz, J.A. Musick, and J. Wyneken (Eds.), The Biology of Sea Turtles Vol. 2. Boca Raton, Florida: crc Press. Pp. 307-338.
- Center for Cetacen Research and Conservation, 2004. Cook Islands Turtle Survey, Palmerston Atoll, 2000. <http://www.whaleresearch.org/turtles/home.htm>. Accessed 9/10/2010.
- Chaloupka, M. and C. Limpus. 2001. Trends in the abundance of sea turtles resident in southern Great Barrier Reef waters. *Biological Conservation* 102:235-249.
- Chaloupka, M., K.A. Bjorndal, G.H. Balazs, A.B. Bolten, L.M. Ehrhart, C.J. Limpus, H. Suganuma, S. Troëng, and M. Yamaguchi. 2007. Encouraging outlook for

- recovery of a once severely exploited marine megaherbivore. *Global Ecology and Biogeography*. DOI: 10.1111/j.1466-8238.2007.00367.x
- Chaloupka, M., G.H. Balazs, S.K.K. Murakawa, R. Morris, and T.M. Work. 2008. Cause-specific spatial and temporal trends in green sea turtle strandings in the Hawaiian Archipelago. *Marine Biology* 154:887-898.
- Craig, P., D. Parker, R. Brainard, M. Rice, and G. Balazs. 2004. Migrations of green turtles in the South Pacific. *Biological Conservation* 116:433-438.
- Crouse, D.T. 1999. The consequences of delayed maturity in a human-dominated world. *American Fisheries Society Symposium* 23: 195-202.
- Cruce, J. 2008. Monitoring of nesting green turtles (*Chelonia mydas*) on Loosiep Island, Ulithi Atoll, Yap, Federated States of Micronesia. Final Contract Report Prepared for JIMAR and NMFS PIRO. 20pp.
- Cruce, J. 2006. Yap State Marine Turtle Conservation Program Ulithi Tagging Project, Gielop and Iar Islands, Summer 2005. Marine Resources Management Division, Colonia, Yap, Federated States of Micronesia. Unpublished Annual Contract Report to NMFS PIRO. 22 pp.
- Cruce, J. 2007. Yap State Marine Turtle Conservation Program Ulithi Tagging Project, Gielop Island Summer 2006. Yap Community Action Program, Colonia, Yap, Federated States of Micronesia. Unpublished Annual Contract Report to NMFS PIRO. 28 pp.
- CNMI Division of Fish and Wildlife (DFW)a. 2009. Population dynamics of sea turtles at the Northern Marianas. Annual Report to NMFS PIRO: October 1, 2008 to September 30, 2009. Grant Number NA08NMF4540613.
- CNMI Division of Fish and Wildlife (DFW)b. Hunting Regulations. <http://www.dfw.gov.mp/Enforcement/Hunting%20Regulations.html>. Accessed 9/10/2010.
- Division of Aquatic and Wildlife Resources (DAWR). 2004. Guam Sea Turtle Recovery Annual Progress Report. March 1, 2004 – August 31, 2004. Grant Progress Report to NMFS PIRO. 9pp.
- Division of Aquatic and Wildlife Resources (DAWR). 2009. Final Annual Progress Report for the Guam Sea Turtle Recovery Project. Award Period 8/1/2006 – 7/31/2008. NOAA Fisheries Grant number NA06NMF4540214. 25 pp.
- Dow, W., Eckert, K.L., Palmer, M. and P. Kramer. 2007. Atlas of Sea Turtle Nesting Habitat for the Wider Caribbean Region. WIDECASST Technical Report No.6 2007. 272pgs.
- Dutton, P.H., G.H. Balazs, R.A. LeRoux, S.K.K. Murakawa, P. Zarate, and L.S. Martinez. 2008. Composition of Hawaiiin green turtle foraging aggregations: mtDNA evidence for a distinct regional population. *Endangered Species Research* 5:37-44.
- Dutton, P.H, and D. Squires. 2008. Reconciling Biodiversity with Fishing: A holistic strategy for Pacific sea turtle recovery. *Ocean Development and International Law*, 39(2): 200-222.
- Dutton, P. 2009 unpublished. Pacific Green Turtle Genetics. A report prepared for the Western Pacific Fishery Management Council's Scientific and Statistical Committee Meeting in Kona, Hawaii, July 22-25, 2009.

- Eckert, K.L. 1993. The biology and population status of marine turtles in the North Pacific. NOAA-TM-NMFS-SWFSC-186.
- Edson, C. and F. Curren. 1987. Report from Oroluk. *Marine Turtle Newsletter* 41:1-2.
- Falanruw, M.V.C. 1971. Island News and Comment. *Atoll Research Bulletin No. 148*. Federated States of Micronesia Code.
- [http://www.fsmlaw.org/fsm/code/title23/T23\\_Ch01.htm](http://www.fsmlaw.org/fsm/code/title23/T23_Ch01.htm). Accessed 9/10/2010.
- Fletcher, M. and G. Petro. 2009. WSB-NOAA Vanuatu Leatherback Monitoring and Outreach Activities: 2008-2009. Wan Smolbag Environment Department unpublished contract report
- Folaumoetu'I, P. 2006. National Biodiversity Strategy and Action Plan. Tonga Department of Environment. Available from: <http://www.sprep.org/att/IRC/eCOPIES/Countries/Tonga/9.pdf>.
- Frazier J (2003) Prehistoric and ancient historic interactions between humans and marine turtles. In: Lutz PL, Musick JA, Wyneken J (eds) *The Biology of sea turtles*, vol 2. CRC Press, Boca Raton, Florida, pp 1–38.
- Geermans, S. 1993. Summary of traditional usage of turtles within the SPREP region. 23pp.
- Government of Fiji. 1992. Fiji Fisheries Act. 25pp.
- Government of Niue 2001. Niue National Biodiversity Strategy and Action Plan. Available from: <http://www.sprep.org/att/IRC/eCOPIES/Contries/Niue/4.pdf>
- Government of Tuvalu. 1975. Wildlife Conservation Ordinance. 12pp.
- Grant, G. S., P. Craig, and G. H. Balazs. 1997. Notes on juvenile hawksbill and green turtles in American Samoa. *Pacific Science* 1:48-53.
- Grimm, G. and J. Farley. 2008. DRAFT Sea turtle nesting activity on Navy Base Guam 2007-2008. NAVFAC Marianas Report. 7 pp.
- Groombridge, B. and R. Luxmoore. 1989. *The Green Turtle and Hawksbill (Reptilia: Cheloniidae): World Status, Exploitation and Trade*. CITES Secretariat, Lausanne, Switzerland. 601 pp.
- Guinea, M. 1993. Sea turtles of Fiji. Report for the South Pacific Regional Environment Program. 52pp.
- Harvey, T., S. Townsend, N. Kenyon, and G. Redfern. 2005. Monitoring of nesting sea turtles in the Coringa Herald National Nature Reserve: 1991/92 to 2003/04 nesting seasons. Indo-Pacific Sea Turtle Conservation Group, report for the Dept. of the Environ. and Heritage. 67pp.
- Havea, S. and K.T. MacKay. 2009. Marine turtle hunting in the Ha`apai Group, Tonga. *Marine Turtle Newsletter* 123: 15-17.
- Hawaii Administrative Rules. Title 13, Subtitle 5, Part 2, Chapter 124. <http://www.state.hi.us/dlnr/dofaw/rules/Chap124.pdf>. accessed 9/10/2010.
- Hawaii State Legislature. [http://www.capitol.hawaii.gov/hrscurrent/Vol03\\_Ch0121-0200D/HRS0195D/HRS\\_0195D-0004.htm](http://www.capitol.hawaii.gov/hrscurrent/Vol03_Ch0121-0200D/HRS0195D/HRS_0195D-0004.htm). Accessed 9/10/2010.
- Heppell, S. S. 1998. Application of life-history theory and population model analysis to turtle conservation. *Copeia*; v. 1998, no. 2, p. 367-375.
- Helfrich, P. 1974. Notes for the ICLARM file on the Cook Islands fisheries organization. November 18, 1974. 1p.
- Hickey, F.R. 2007. Traditional marine resource management in Vanuatu; world views in transformation. p147-168 In: Haggan, N., B. Neis, and I.G. Baird (eds.). *Fishers*

- Knowledge in Fisheries Science and Management. Coastal Management Sourcebooks 4. Paris: UNESCO, 437 pp.
- Hirth, H.F. 1993. Nearshore Marine Resources of the South Pacific: Chapter 10, Marine Turtles. Wright, A. and L. Hill (eds.). Honiara, Suva: Forum Fisheries Agency, Institute of Pacific Studies.
- Hirth, H. F. 1997. Synopsis of biological data on the green turtle *Chelonia mydas*. U.S. Dept. of the Interior, Fish and Wildlife Service, Biological Report 97(1); 1997, 120 pp.
- Ilo, L., G.P. Camacho, and C. Alepuyo. 2005. 2005 Sea turtle nesting and in-water assessment report for the Commonwealth of the Northern Mariana southern inhabitant islands of Saipan, Tinian, and Rota. CNMI Division of Fish and Wildlife Report. 23 pp.
- Ilo, L. and J. Mangolona. 2002. Sea Turtle Assessment Report for the Island Anatahan. Sept. 27, 2002. CNMI DFW unpublished report.
- Ishizaki, A., 2007. Towards Reducing Human-caused Impacts on Green Turtle (*Chelonia mydas*) Nesting Activity in Ogasawara Islands, Japan. Thesis, Colorado State University.
- Johannes, R.E. 1978. Traditional marine conservation methods in Oceania and their demise. *Annual Review of Ecology and Systematics* 9:349-364.
- Johannes, R.E. 1986. A review of information on the subsistence use of green and hawksbill sea turtles on islands under United States jurisdiction in the Western Pacific Ocean. US Department of Commerce National Oceanographic and Atmospheric Administration National Marine Fisheries Service Administrative Report SWR-86-2. 41: 28-29.
- Kamezaki, N., Kikukawa, A., and H. Ota. 1999. Current status of the sea turtles nesting on Okinawajima and adjacent islands of the central Ryukus, Japan. *Biological Conservation*, vol. 87(1): 149-153.
- Kinan, I. and P. Dalzel. 2004. Turtle conservation and fisheries development in the Pacific Islands: Different perspectives create conflicts between developed and developing economies. Sea turtles: Flagship species for conservation and fishery management. *Maritime Studies*. 3(2): 195-212.
- Kinch, J. 2003a. Sea Turtle Resources in the Milne Bay Province, Papua New Guinea: Results of a Nesting Survey (21-27/01/03) at Panayayapona and Panadaludalu Islands (Jomard Islands), with Additional Notes. Milne Bay Community-based Coastal and Marine Conservation Program unpublished report.
- Kinch, J. 2003b. Sea turtle resources in the Milne Bay Province, Papua New Guinea: Conservation and management issues in relation to protection and sale in markets. Milne Bay Community-based Coastal and Marine Conservation Program unpublished report.
- Kinch, J. 2006. Socioeconomic baseline study of communities involved in leatherback turtle nesting beach projects along the Huon Coast, Morobe Province, Papua New Guinea. Final report prepared for the Western Pacific Regional Fishery Management Council, Honolulu, Hawaii.  
[http://www.wpcouncil.org/protected/Documents/Kinch%20report\\_final.pdf](http://www.wpcouncil.org/protected/Documents/Kinch%20report_final.pdf)

- Kolinski, S. 1992. Outer Islands Turtle Project: Stage II; Report on the Gielop Island Fieldwork. Report prepared for the Marine Resources Management Division, Yap State, Federated States of Micronesia.
- Kolinski, S. 1993. Outer Island Turtle Project: Stage III; Report on Elato Atoll Fieldwork. Report prepared for the Marine Resources Management Division, Yap State, Federated States of Micronesia.
- Kolinski, S.P., D.M. Parker, L.I. Ilo, and J.K. Ruak. 2001. An assessment of the sea turtles and their marine and terrestrial habitats at Saipan, Commonwealth of the Northern Mariana Islands. *Micronesica* 34(1):55-72.
- Kolinski, S, L.I. Ilo and J.M. Manglona. 2004. Green turtles and their marine habitats at Tinian and Aguijan, with projections on resident turtle demographics in the southern arc of the Commonwealth of the Northern Mariana Islands. *Micronesica* 37:95-116.
- Kolinski, S.P., Cruce, J., Parker, D.M., Frutchey, K.P., Balazs, G.H., Clarke, R. 2007 unpublished manuscript. Identifying Migration-Based Connectivity via Satellite Telemetry for Post-Nesting Green Turtles from Gielop Island, Federated States of Micronesia
- Laveti, M. and K.T. MacKay. 2009. Does Fiji's turtle moratorium work? *Marine Turtle Newsletter* 123:12-15.
- Leary, T. and M. Laumani. 1989. Marine turtles of Isabel Province: A report of a survey of nesting beaches (7-21 November 1989). Environment and Conservation Division and Fisheries Division MNR. 14pp.
- Lebeau, A. 1985. Breeding evaluation trials in the green turtle *Chelonia mydas* (Linne) on Scilly Atoll (Leeward Islands, French Polynesia) during the breeding season 1982-1983 and 1983-1984. Proceedings of the Fifth International Coral Reef Congress, Tahiti, 5:487-493.
- Lessa, W. 1983. Sea turtles and ritual: conservations in the Caroline Islands. In, The fishing culture of the world, B. Gunda (ed.). Akademiai Kiado, publishing house of the Hungarian Academy of Sciences. Budapest, pages 1183-1201.
- Limpus, C.J., J.D. Miller, C.J. Parmenter, and D.J. Limpus. 2003. The Green Turtle, *Chelonia mydas*, population of Raine Island and the northern Great Barrier Reef: 1843-2001. *Memoirs of the Queensland Museum* 49(1):349-440. Brisbane. ISSN 0079-8835.
- Limpus, C.J. 2009. A biological review of Australian marine turtle species. The State of Queensland, Environmental Protection Agency.
- Limpus, C., S. Mounier, and A. Downer. 2009 (in prep). Project to determine the distribution and population sizes of nesting marine turtles in New Caledonia, and to enhance the capacity of local governments to protect and manage these species. Poster presented in the Twenty-ninth Annual Sea Turtle Symposium in Brisbane Australia, February, 2009.
- McClenachan, L., Jackson, J. B. C.; and M.J.H. Newman. 2006. Conservation implications of historic sea turtle nesting beach loss. *Frontiers in Ecology and the Environment*; 4 (6): 290-296.
- McCoy, M.A. 1974. Man and Turtle in the Central Carolines. *Micronesica* 10(2): 207-221.



- McCoy, M.A. 1997. The traditional and ceremonial use of the Green sea turtle (*Chelonia mydas*) in the Northern Mariana Islands: With recommendations for its use in cultural events and education. A report prepared for the Western Pacific Regional Fishery Management Council & University of Hawai'i, Sea Grant College Program.
- McCoy, M. 2004. Defining parameters for sea turtle research in the Marshall Islands. NOAA ADMIN REPORT AR-PIR-08-04.
- McKeown, A. 1977. Marine turtles of the Solomon Islands. Ministry of Natural Resources, Fisheries Division. 49 pp.
- Melyan, A.B. 1982. Sea turtle migration-evidence from tag returns. In; KA. Bjorndal (ed), *Biology and Conservation of Sea Turtles*. Smithsonian Institution Press, Washington DC. Pp. 91-100.
- Miller DJ. 1997. Reproduction in sea turtles. In *The Biology of Sea Turtles*, Lutz PL, Musick JA (eds). CRC Publishing: Boca Raton; 51–81.
- Moritz, C., D. Broderick, K. Dethmers, N. Fitzsimmons, and C. Limpus. 2002. Population genetics of Southeast Asian and Western Pacific green turtles, *Chelonia mydas*. Final Report to UNEP/CMS, 20 June 2002.
- National Fisheries Authority (NFA). 2007. A review of fisheries and marine resources in New Ireland province, Papua New Guinea. National Fisheries Authority and Coastal Fisheries Management and Development Project, 2007. 47 pp.
- National Marine Fisheries Service and U.S. Fish and Wildlife Service. 1998. Recovery Plan for U.S. Pacific Populations of the Green Turtle. 84pp.
- National Marine Fisheries Service and U.S. Fish & Wildlife Service. 2007. Green Sea Turtle (*Chelonia mydas*). 5-Year Review: Summary and Evaluation. 105pp.
- National Research Council (NRC). 2010. Assessment of sea turtle status and trends: Integrating demography and abundance. Committee on the review of sea turtle population assessment methods. The National Academy of Sciences, National Academies Press, Washington, DC. 143 pp.
- Naughton, J.J. 2001. Sea turtle survey at Oroluk Atoll and Minto Reef, Federated States of Micronesia. *Marine Turtle Newsletter* 55:9-12.
- Neithammer, K.R., Balazs, G.H., Hatfield, J.S., Nakai, G.L. and J.L. Megysi. 1997. Reproductive Biology of the Green Turtle (*Chelonia mydas*) at Tern Island, French Frigate Shoals, Hawai'i. *Pacific Science*, vol. 51: 1: 36-47
- PacLII. Laws of the Republic of Kiribati Revised Edition 1979 Chapter 33 Fisheries. [http://www.pacii.org/ki/legis/consol\\_act/fo163/](http://www.pacii.org/ki/legis/consol_act/fo163/). Accessed 9/10/2010.
- Palau Bureau of Marine Resources. 2005. Palau Marine Turtle Conservation & Monitoring Program Final Report. Unpublished grant report. 70pp.
- Palau Bureau of Marine Resources. 2008. Palau Marine Turtle Conservation & Monitoring Program Final Report. Unpublished grant report. 31pp.
- Petro, G., F. Hickey, and K. MacKay. 2007. Leatherback turtles in Vanuatu. *Chelonian Conservation and Biology* 6(1):135-137.
- Pita, E. 1979. The turtle status in Tuvalu. Joint SPC-NMFS Workshop on Marine Turtles in the Tropical Pacific Islands. Noumea, New Caledonia 11-14 December, 1979. 3pp.
- Pita, J. and D. Broderick. 2005. Hawksbill turtles in the Solomon Islands. IN: I. Kinan (ed.) *Proceedings of the Second Western Pacific Sea Turtle Cooperative Research*

- & Management Workshop. Western Pacific Regional Fisheries Management Council, Honolulu, Hawaii, May 17-21, 2004.
- Pritchard, P.C.H. 1977. Marine Turtles of Micronesia. Chelonia Press, San Francisco, California. 83pp.
- Pritchard, P.C.H. 1979. Marine turtles of Papua New Guinea. Unedited field notes. Account of fieldwork conducted on behalf of Papua New Guinea Wildlife Division by Peter C.H. Pritchard and Suzanne Rayner, August to October 1978. Fisheries Archive Paper No. P87. 63pp.
- Pritchard, P.C.H. 1982. Nesting of the leatherback turtle in Pacific Mexico with a new estimate of the world population status. *Copeia* 1982(4):741-747.
- Pritchard, P.C.H. 1994. Les D'Entrecasteaux Enfin! Report of an expedition to study the sea turtles of the D'Entrecasteaux reefs, north of New Caledonia. In: Schroeder, B.A., and B.E. Witherington (compilers). Proceedings of the Thirteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-341:143-145.
- Pritchard, P.C.H. 1995a. Marine turtles of the south Pacific. In: Bjorndal, K. (ed.) Biology and Conservation of Sea Turtles. Smithsonian Institution Press. Washington D.C., 615pp.
- Pritchard, P.C.H. 1995b. Marine turtles of Micronesia. In: Bjorndal, K. (ed.) Biology and Conservation of Sea Turtles. Smithsonian Institution Press. Washington D.C., 615pp.
- Pulea, M. 1992. Legislative Review of environmental Law, Cook Islands. SPREP Regional Tech Assistance Project II Titel III (Series). Available from: [http://www.sprep.org/att/IRC/eCOPIES/Countries/Cook\\_Islands/10.pdf](http://www.sprep.org/att/IRC/eCOPIES/Countries/Cook_Islands/10.pdf).
- Puleloa, W.K. and N. Kilma. 1992. The Sea Turtles of the Northern Marshalls: A research expedition to Bikar and Erikup Atolls, and Jemo Island. Unpublished Report, 72pp.
- Pultz, S., D. O'Daniel, S. Krueger, H. McSharry, and G. Balazs. 1999. Marine turtle survey on Tinian, Mariana Islands. *Micronesica* 31(2):85-94.
- Ramohia, P.C. and J. Pita. 1996. Arnavons Islands Survey: Report to SPREP on the Regional Marine Turtle Conservation Programme (1995 project) in the Solomon Islands. Fisheries Division of the Ministry of Agriculture and Fisheries, Honiara. 19pp.
- Republic of the Marshall Islands. 1997. Marine Resources Act. 70pp.
- Schroeder, B. and S. Murphy. 1999. "Population surveys (ground and aerial) on nesting beaches." pp. 45-53 In: K.L. Eckert, K.A. Bjorndal, F.A. Abreu-Grobis and M. Donnelly (eds.). *Research and Management Techniques for the Conservation of Sea Turtles*. IUCN/SSC Marine Turtle Specialist Group.
- Secretariat of the Pacific Regional Environment Program (SPREP). 2007. Summary of in-country legislation. SPREP unpublished report, Apia Samoa.
- Secretariat of the Pacific Regional Environment Program (SPREP). 2008. 2008-2012 Marine Turtle Action Plan. SPREP technical report, Apia Samoa.
- Semioff, J.A. 2004. 2004 Global Status Assessment: Green turtle (*Chelonia mydas*). IUCN Marine Turtle Specialist Group Review. 71pp.
- Smith, A., S. Kolinski and V. Hachiglou. 1991. Outer Island Turtle Project, Yap State, F.S.M. – Status Report. Report prepared for the South Pacific Regional

- Environment Programme's Regional Marine Turtle Conservation Programme meeting, August 12 & 13 1991, Noumea, New Caledonia. 6pp.
- South Pacific Commission. 1979a. Notes on marine turtles of Republic of Kiribati. Joint SPC-NMFS Workshop on marine turtles in the tropical Pacific Islands. Noumea, New Caledonia, 11-14 December, 1979. 16pp.
- South Pacific Commission. 1979b. Tagging and rearing of the green turtle *Chelonia mydas* conducted in French Polynesia by the Department of Fisheries. Joint SPC-NMFS Workshop on Marine Turtles in the Tropical Pacific Islands. Noumea, New Caledonia, 11-14 December, 1979. 22pp.
- South Pacific Commission and Bureau of Marine Resources Palau. 2007. Palau Domestic Fishing Laws. 52pp.
- South Pacific Projects. 2010. Marine Turtle Steering Committee. <http://www.southpacificprojects.org/marine-turtle-steering-committee/>. Accessed 9/10/2010.
- Spring, C. S. 1982. Status of marine turtle populations in Papua New Guinea. In: Bjorndal, K. (ed.) *Biology and Conservation of Sea Turtles*. Smithsonian Institution Press. Washington D.C., 615pp.
- Stone, G., D. Obura, S. Bailey, A. Yoshinaga, C. Holloway, R. Barrel, and S. Mangubhai. 2001. *Marine Biological Surveys of the Phoenix Islands: Summary of Expedition Conducted from June 24-July 15, 2000*. New England Aquarium. 107pp.
- Tagarino, A., K.S. Saili, and R. Utzurrum. 2008. Investigations into the Status of Marine Turtles in American Samoa, with Remediation of Identified Threats and Impediments to Conservation and Recovery of Species. NOAA Grant Award No. NA04NMF4540126. FINAL REPORT (01 October 2004 to 30 September 2008), 44pp.
- Tagarino, A. and R. Utzurrum. 2010. Investigations into the Status of Marine Turtles in American Samoa: Assessment of Threat to Nesting Activities and Habitat in Swains Island. NOAA Grant Award No. NA08NMF4540506. FINAL REPORT (01 October 2008 to 30 September 2009), 14pp.
- Te Mana o Te Moana. 2009. Marine Environment Protection and Public Awareness in French Polynesia: Te Mana o te Moana Activity Report Since 4 Years 2004-2008. 4pp. <http://www.temanaotemoana.org/downloads/activity-report.pdf>.
- Te mana Tea. 2007. Project Tikehau. Contract unpublished report. In French.
- Troeng, S., and M. Chaloupka. 2007. Variation in adult annual survival probability and remigration intervals of sea turtles. *Marine Biology* 151(5):1721-1730.
- Tuato'o-Bartley, N., T.E. Morrell, and P. Craig. 1993. Status of Sea Turtles in American Samoa in 1991. *Pacific Science* 47(3):215-221.
- United States Navy (USN). 2010. Biological Assessment: Carrier vessel nuclear (CVN) warf construction and Navy waterfront improvement, Apra Harbor, Guam. NAVFAC PAC March 12, 2010.
- Utzurrum, R. 2002. Sea turtle conservation in American Samoa. In: Kinan, I. (ed.). *Proceedings of the Western Pacific Sea Turtle Cooperative Research and Management Workshop*. February 5-8, 2002, Honolulu, Hawaii, USA. Western Pacific Regional Fishery Management Council. 300pp.
- Van Buskirk, J. and L.B. Crowder. 1994. Life-history variation in marine turtles. *Copeia*; v. 1994: 66-81.

- Vaughan, P. 1980. Marine Turtles of Santa Isabel Province: survey report. Fisheries Division, Ministry of Natural Resources technical report.
- Vaughan, P. 1981. Marine turtles: A review of their status and management in the Solomon Islands. Fisheries Division, Ministry of Natural Resources technical report.
- Vogt, S. 2009. Wildlife Surveys on Military Leased Lands, Farallon de Medinilla, Commonwealth of the Northern Mariana Islands: Fiscal Years 2006-2007. Annual Navy NAVFAC Report for 61755NR410.
- Wanguu, N., D. Kwan, I. Bell, and J. Pita. 2004. Turtle tagging and monitoring in Milne Bay Province, December 2003. A report prepared for Conservation International, Papua New Guinea Department of Environment and Conservation and South Pacific Regional Environmental Program. Milne Bay Community-based Marine Conservation Program. PNG 01/010. 45pp.
- Weaver, S. 1996. The Fiji Sea Turtle Conservation Strategy. WWF Project No. 9p005.01. 54pp.
- Witzell, W.N. 1982. Observations on the green sea turtle (*Chelonia mydas*) in Western Samoa. *Copeia* 1:183-185.
- Woodrom-Luna, R. 2003. The merging of archaeological evidence and marine turtle ecology: A case study approach to the importance of including archaeological data in marine science. *SPC Traditional Marine Resource Management and Knowledge Information Bulletin*; 15: 26-30.
- Woodrom-Rudrud, R., J. Walsh Kroeker, H. Young Leslie and S. Finney. 2007. Sea turtles wars: Culture, war and sea turtles in the Republic of the Marshall Islands. *SPC Traditional Marine Resource Management and Knowledge Information Bulletin*; 21: 3-29.
- Woodrum-Rudrud, R. 2010. Forbidden Sea Turtles: Traditional laws pertaining to sea turtle consumption in Polynesia (including the Polynesian outliers). *Conservation and Society*; 8(1): 84-97.
- Yap State Code, Title 18, Division 3, Chapter 10, § 1005.